

GERSTENHABER STRUCTURE ON HOCHSCHILD COHOMOLOGY  
OF THE FOMIN-KIRILLOV ALGEBRA ON 3 GENERATORS

ESTANISLAO HERSCOVICH AND ZILING LI

Received: April 11, 2022

Revised: September 9, 2022

Communicated by Henning Krause

ABSTRACT. The goal of this article is to compute the Gerstenhaber bracket of the Hochschild cohomology of the Fomin-Kirillov algebra on three generators over a field of characteristic different from 2 and 3. This is in part based on a general method we introduce to easily compute the Gerstenhaber bracket between elements of  $\mathrm{HH}^0(A)$  and elements of  $\mathrm{HH}^n(A)$  for  $n \in \mathbb{N}_0$ , the method by M. Suárez-Álvarez in [8] to calculate the Gerstenhaber bracket between elements of  $\mathrm{HH}^1(A)$  and elements of  $\mathrm{HH}^n(A)$  for any  $n \in \mathbb{N}_0$ , as well as an elementary result that allows to compute the remaining brackets from the previous ones. We also show that the Gerstenhaber bracket of  $\mathrm{HH}^\bullet(A)$  is not induced by any Batalin-Vilkovisky generator.

2020 Mathematics Subject Classification: 16E40, 18G10, 16S37

Keywords and Phrases: Fomin-Kirillov algebra, Hochschild cohomology, Gerstenhaber bracket

## 1 INTRODUCTION

The goal of this article is to completely compute the Gerstenhaber bracket of the Hochschild cohomology  $\mathrm{HH}^\bullet(A)$  of the Fomin-Kirillov algebra  $A$  on three generators over a field  $\mathbb{k}$  of characteristic different from 2 and 3. In the paper [3] we have computed the algebra structure on  $\mathrm{HH}^\bullet(A)$ , and we showed that it is a finitely generated graded algebra with a minimal set  $\{X_i | i \in [1, 14]\}$  of 14 generators, three of which are in  $\mathrm{HH}^0(A)$ , five are in  $\mathrm{HH}^1(A)$ , four are in  $\mathrm{HH}^2(A)$ , one in  $\mathrm{HH}^3(A)$  and one in  $\mathrm{HH}^4(A)$ . In this article we compute the Gerstenhaber bracket between these generators, which fully determines the Gerstenhaber bracket on  $\mathrm{HH}^\bullet(A)$ .

Our calculations are organised as follows. First we provide a general method of homological flavour to easily compute the Gerstenhaber bracket between elements of  $\mathrm{HH}^0(A)$  and elements of  $\mathrm{HH}^n(A)$  for any  $n \in \mathbb{N}_0$  and any algebra  $A$  over a field  $\mathbb{k}$  (see Theorem 2.7). We then use this method for the particular case of the Fomin-Kirillov algebra  $A$  on three generators over a field of characteristic different from 2 and 3 (see Proposition 4.4). Secondly, for  $A$  as before, we compute the Gerstenhaber bracket between elements of  $\mathrm{HH}^1(A)$  and elements of  $\mathrm{HH}^n(A)$  for any  $n \in \mathbb{N}_0$  using the method introduced by M. Suárez-Álvarez in [8] (see Propositions 4.5 and 4.10). Finally, we present a simple result that allows us to compute the remaining Gerstenhaber brackets under some assumptions on the algebra structure of the Hochschild cohomology of an algebra (see Lemma 4.12), which are verified in the case of the Fomin-Kirillov algebra  $A$  on three generators over a field of characteristic different from 2 and 3 (see Proposition 4.14). We summarize our results in Table 4.1. By using the explicit expression of the Gerstenhaber bracket of  $\mathrm{HH}^\bullet(A)$  together with a direct computation, we show that the former is not induced by any Batalin-Vilkovisky generator (see Proposition 4.15).

The article is organised as follows. In the first subsection of Section 2 we recall some general facts about the Gerstenhaber bracket on Hochschild cohomology. In Subsection 2.2 we present our general method to compute the Gerstenhaber bracket between elements of  $\mathrm{HH}^0(A)$  and elements of  $\mathrm{HH}^n(A)$  for any  $n \in \mathbb{N}_0$ , whereas in Subsection 2.3 we recall the method of [8] to compute the Gerstenhaber bracket between elements of  $\mathrm{HH}^1(A)$  and elements of  $\mathrm{HH}^n(A)$  for any  $n \in \mathbb{N}_0$ . In Section 3 we recall the definition of the Fomin-Kirillov algebra on three generators, and we summarize some results of [3] about its Hochschild cohomology, in particular those we are going to use in the sequel. Finally, in Section 4 we compute the Gerstenhaber brackets of the generators of the algebra structure of the Hochschild cohomology of the Fomin-Kirillov algebra  $A$  on three generators, following the path described in the previous paragraph.

In the whole article, we will denote by  $\mathbb{N}$  (resp.,  $\mathbb{N}_0$ ) the set of positive (resp., nonnegative) integers, and  $\mathbb{Z}$  the set of integers. Moreover, given  $i, j \in \mathbb{Z}$  we define the integer interval  $\llbracket i, j \rrbracket = \{n \in \mathbb{Z} \mid i \leq n \leq j\}$ . To reduce space in the expressions of the article we will typically denote the composition  $f \circ g$  of maps  $f$  and  $g$  simply by their juxtaposition  $fg$ . For a field  $\mathbb{k}$ , all maps between  $\mathbb{k}$ -vector spaces will be  $\mathbb{k}$ -linear and all unadorned tensor products  $\otimes$  will be over  $\mathbb{k}$ .

We would like to thank the referee for the careful reading of the article.

## 2 THE GERSTENHABER BRACKET ON HOCHSCHILD COHOMOLOGY

All along this section we will consider  $\mathbb{k}$  to be a field and  $A$  to be a (unital associative)  $\mathbb{k}$ -algebra.

2.1 GENERALITIES ON THE GERSTENHABER BRACKET

In this subsection we recall several basic definitions and results concerning the Gerstenhaber bracket, that we will utilize in the sequel.

Recall that the **BAR RESOLUTION**  $(B_\bullet(A), d_\bullet)$  of  $A$  is given by  $B_n(A) = A^{\otimes(n+2)}$  for  $n \in \mathbb{N}_0$ , with the differentials  $d_n : B_n(A) \rightarrow B_{n-1}(A)$  given by

$$d_n(a_0 | \dots | a_{n+1}) = \sum_{j=0}^n (-1)^j a_0 | \dots | a_{j-1} | a_j a_{j+1} | a_{j+2} | \dots | a_{n+1}$$

for  $a_0, \dots, a_{n+1} \in A$  and  $n \in \mathbb{N}$ , and the augmentation  $\pi : B_0(A) = A \otimes A \rightarrow A$  defined by the multiplication of  $A$ . We will typically write  $a_0 | \dots | a_{n+1}$  instead of  $a_0 \otimes \dots \otimes a_{n+1}$  for simplicity. There is an isomorphism

$$F : \mathrm{Hom}_{A^e}(B_n(A), A) \longrightarrow \mathrm{Hom}_{\mathbb{k}}(A^{\otimes n}, A)$$

given by  $F(f)(a_1 | \dots | a_n) = f(1 | a_1 | \dots | a_n | 1)$  for  $f \in \mathrm{Hom}_{A^e}(B_n(A), A)$  and  $a_1, \dots, a_n \in A$ . The inverse map

$$G : \mathrm{Hom}_{\mathbb{k}}(A^{\otimes n}, A) \longrightarrow \mathrm{Hom}_{A^e}(B_n(A), A)$$

of  $F$  is explicitly given by  $G(g)(a_0 | \dots | a_{n+1}) = a_0 g(a_1 | \dots | a_n) a_{n+1}$  for  $g \in \mathrm{Hom}_{\mathbb{k}}(A^{\otimes n}, A)$  and  $a_0, \dots, a_{n+1} \in A$ .

The following definition is classical (see for instance [10], Def. 1.4.1).

**DEFINITION 2.1.** *Let  $m, n \in \mathbb{N}_0$ ,  $f \in \mathrm{Hom}_{\mathbb{k}}(A^{\otimes m}, A)$  and  $g \in \mathrm{Hom}_{\mathbb{k}}(A^{\otimes n}, A)$ . The **GERSTENHABER BRACKET**  $[f, g]$  is defined at the chain level as the element of  $\mathrm{Hom}_{\mathbb{k}}(A^{\otimes(m+n-1)}, A)$  given by*

$$[f, g] = f \circ_G g - (-1)^{(m-1)(n-1)} g \circ_G f,$$

where  $f \circ_G g$  is defined by

$$\begin{aligned} & (f \circ_G g)(a_1 | \dots | a_{m+n-1}) \\ &= \sum_{i=1}^m (-1)^{(n-1)(i-1)} f(a_1 | \dots | a_{i-1} | g(a_i | \dots | a_{i+n-1}) | a_{i+n} | \dots | a_{m+n-1}). \end{aligned}$$

Moreover, if  $m = 0$ , then  $f \circ_G g = 0$ , while if  $n = 0$ , then the formula should be interpreted by taking the value  $g(1)$  in place of  $g(a_i | \dots | a_{i+n-1})$ .

Using the isomorphisms  $F$  and  $G$  of chain complexes given above, one defines the **GERSTENHABER BRACKET** in  $\mathrm{Hom}_{A^e}(B_\bullet(A), A)$  by  $[f, g] = G([F(f), F(g)]) \in \mathrm{Hom}_{A^e}(B_{m+n-1}(A), A)$  for  $f \in \mathrm{Hom}_{A^e}(B_m(A), A)$ ,  $g \in \mathrm{Hom}_{A^e}(B_n(A), A)$  and  $m, n \in \mathbb{N}_0$ . The Gerstenhaber bracket given before induces a well-defined bilinear map

$$\begin{aligned} [\cdot, \cdot] : \mathrm{H}^m(\mathrm{Hom}_{A^e}(B_\bullet(A), A)) \times \mathrm{H}^n(\mathrm{Hom}_{A^e}(B_\bullet(A), A)) \\ \longrightarrow \mathrm{H}^{m+n-1}(\mathrm{Hom}_{A^e}(B_\bullet(A), A)) \end{aligned}$$

for all  $m, n \in \mathbb{N}_0$ , that we also call the GERSTENHABER BRACKET. More generally, let  $(P_\bullet, \partial_\bullet)$  be a projective bimodule resolution over  $A$  with augmentation  $\mu : P_0 \rightarrow A$ . Let  $i_\bullet : P_\bullet \rightarrow B_\bullet(A)$  and  $p_\bullet : B_\bullet(A) \rightarrow P_\bullet$  be morphisms of complexes of  $A$ -bimodules lifting  $\text{id}_A$ , so  $p_\bullet i_\bullet$  is homotopic to  $\text{id}_{P_\bullet}$  and  $i_\bullet p_\bullet$  is homotopic to  $\text{id}_{B_\bullet(A)}$ . We also recall that the morphisms  $i_\bullet$  and  $p_\bullet$  induce the quasi-isomorphisms  $i_\bullet^* : \text{Hom}_{A^e}(B_\bullet(A), A) \rightarrow \text{Hom}_{A^e}(P_\bullet, A)$  and  $p_\bullet^* : \text{Hom}_{A^e}(P_\bullet, A) \rightarrow \text{Hom}_{A^e}(B_\bullet(A), A)$  given by  $i_\bullet^*(f) = f i_\bullet$  and  $p_\bullet^*(g) = g p_\bullet$  for  $f \in \text{Hom}_{A^e}(B_\bullet(A), A)$  and  $g \in \text{Hom}_{A^e}(P_\bullet, A)$ , respectively. Moreover,  $H(i_\bullet^*), H(p_\bullet^*) : \text{HH}^\bullet(A) \rightarrow \text{HH}^\bullet(A)$  are independent of the choice of  $i_\bullet$  and  $p_\bullet$ . The GERSTENHABER BRACKET

$$[\cdot, \cdot] : H^m(\text{Hom}_{A^e}(P_\bullet(A), A)) \times H^n(\text{Hom}_{A^e}(P_\bullet(A), A)) \longrightarrow H^{m+n-1}(\text{Hom}_{A^e}(P_\bullet(A), A))$$

for all  $m, n \in \mathbb{N}_0$  is then defined by transport of structures. More generally, given cocycles  $f \in \text{Hom}_{A^e}(P_m, A)$  and  $g \in \text{Hom}_{A^e}(P_n, A)$ , we define the Gerstenhaber bracket  $[f, g] \in H^{m+n-1}(\text{Hom}_{A^e}(P_\bullet, A)) \cong \text{HH}^{m+n-1}(A)$  as the cohomology class of  $i_\bullet^*([p_\bullet^*(f), p_\bullet^*(g)])$ .

The following properties of the Gerstenhaber bracket are classical (see for instance [2], equation (2), cf. [10], Lemmas 1.4.3 and 1.4.7).

LEMMA 2.2. *Let  $\mathbb{k}$  be a field and  $A$  a  $\mathbb{k}$ -algebra. Then*

$$\begin{aligned} [x, y] &= -(-1)^{(m-1)(n-1)}[y, x], \\ [x, [y, z]] &= [[x, y], z] + (-1)^{(m-1)(n-1)}[y, [x, z]], \end{aligned} \tag{2.1}$$

and

$$[x \smile y, z] = [x, z] \smile y + (-1)^{m(p-1)}x \smile [y, z] \tag{2.2}$$

for all  $x \in \text{HH}^m(A)$ ,  $y \in \text{HH}^n(A)$  and  $z \in \text{HH}^p(A)$ , where  $\smile$  denotes the cup product.

REMARK 2.3. *To avoid any confusion, we remark that the definition of cup product on Hochschild cohomology in the previous result is the one given in [2], Section 7. Explicitly, at the level of the bar resolution it is given by  $(f \smile g)(a_1, \dots, a_{m+n}) = f(a_1, \dots, a_m)g(a_{m+1}, \dots, a_{m+n})$ , for  $a_1, \dots, a_{m+n} \in A$ ,  $f \in \text{Hom}_{\mathbb{k}}(A^{\otimes m}, A)$  and  $g \in \text{Hom}_{\mathbb{k}}(A^{\otimes n}, A)$ . A different convention, in the spirit of Koszul’s sign rule, includes a sign  $(-1)^{mn}$  (see [10], Def. 1.3.1 and Rk. 1.3.3). To reduce space we will denote the cup product simply by juxtaposition.*

The previous result is typically rephrased by stating that the Hochschild cohomology is a GERSTENHABER ALGEBRA, i.e. a graded-commutative algebra  $H = \bigoplus_{n \in \mathbb{N}_0} H^n$  endowed with a bracket  $[\cdot, \cdot] : H \otimes H \rightarrow H$  satisfying  $[H^m, H^n] \subseteq H^{m+n-1}$  for  $m, n \in \mathbb{N}_0$ , (2.1) and (2.2).

Assume for the rest of this subsection that  $A$  is **GRADED**, *i.e.* there exist  $\mathbb{k}$ -vector subspaces  $\{A_n\}_{n \in \mathbb{Z}}$  of  $A$  such that  $A = \bigoplus_{n \in \mathbb{Z}} A_n$ , and  $A_m \cdot A_n \subseteq A_{m+n}$  for all  $m, n \in \mathbb{Z}$ . Recall that a left  $A$ -module  $M$  is called **GRADED** if there are  $\mathbb{k}$ -vector subspaces  $\{M_n\}_{n \in \mathbb{Z}}$  of  $M$  such that  $M = \bigoplus_{n \in \mathbb{Z}} M_n$  and  $A_m \cdot M_n \subseteq M_{m+n}$  for all  $m, n \in \mathbb{Z}$ . Given two graded left  $A$ -modules  $M$  and  $N$ , a morphism  $f : M \rightarrow N$  of left  $A$ -modules is called **HOMOGENEOUS** of degree  $d \in \mathbb{Z}$  if  $f(M_n) \subseteq N_{n+d}$  for all  $n \in \mathbb{Z}$ . Let  $\mathrm{Hom}_A(M, N)$  be the  $\mathbb{k}$ -vector space consisting of all morphisms of left  $A$ -modules from  $M$  to  $N$ . Let

$$\mathcal{H}om_A(M, N) = \bigoplus_{d \in \mathbb{Z}} \mathcal{H}om_A(M, N)_d,$$

be the graded  $\mathbb{k}$ -vector space, where  $\mathcal{H}om_A(M, N)_d$  is the subspace of  $\mathrm{Hom}_A(M, N)$  consisting of all homogeneous morphisms of degree  $d$ . The following result is classical (see [5], Cor. 2.4.4).

LEMMA 2.4. *If  $M$  is a finitely generated graded module over a graded algebra  $A$ , then  $\mathrm{Hom}_A(M, N) = \mathcal{H}om_A(M, N)$ .*

COROLLARY 2.5. *Let  $(P_\bullet, \partial_\bullet)$*

$$\dots \xrightarrow{\partial_3} P_2 \xrightarrow{\partial_2} P_1 \xrightarrow{\partial_1} P_0 \xrightarrow{\mu} A \longrightarrow 0$$

*be a projective bimodule resolution of a graded  $\mathbb{k}$ -algebra  $A$ , where  $P_i$  is finitely generated as left  $A^e$ -module for  $i \in \mathbb{N}_0$ , and  $\mu$  and  $\partial_i$  are homogeneous of degree 0 for  $i \in \mathbb{N}$ . Then  $\mathrm{Hom}_{A^e}(P_i, A) = \mathcal{H}om_{A^e}(P_i, A)$  for  $i \in \mathbb{N}_0$ . Hence, the Hochschild cohomology  $\mathrm{HH}^\bullet(A) \cong \bigoplus_{i \in \mathbb{N}_0} \mathrm{H}^i(\mathrm{Hom}_{A^e}(P_\bullet, A))$  of  $A$  is a bigraded algebra, for the cohomological degree  $i$  and the internal degree induced by that of  $A$  and  $P_\bullet$ . Moreover, the cup product and the Gerstenhaber bracket on  $\mathrm{HH}^\bullet(A)$  preserve the internal degree.*

REMARK 2.6. *The existence of a projective bimodule resolution of the graded  $\mathbb{k}$ -algebra  $A$  satisfying the conditions of the previous corollary clearly holds if the graded  $\mathbb{k}$ -algebra  $A^e$  is noetherian (e.g. if  $A$  is finite dimensional over  $\mathbb{k}$ ).*

## 2.2 METHOD COMPUTING THE BRACKET BETWEEN $\mathrm{HH}^0(A)$ AND $\mathrm{HH}^n(A)$

In this subsection we introduce an elementary method to compute the Gerstenhaber bracket between the cohomology groups  $\mathrm{HH}^0(A)$  and  $\mathrm{HH}^n(A)$  for  $n \in \mathbb{N}_0$  of any algebra  $A$  using any projective bimodule resolution of  $A$  (see Thm. 2.7). We were unable to explicitly find this method in the existing literature (see Remark 2.12), although we suspect it could be well known to the experts.

Let  $\rho$  be an element of the center  $\mathcal{Z}(A) \cong \mathrm{HH}^0(A)$  of  $A$  and  $\ell_\rho \in \mathrm{Hom}_{A^e}(B_0(A), A)$  be the morphism defined by  $\ell_\rho(1|1) = \rho$ . Let  $(P_\bullet, \partial_\bullet)$  be a projective bimodule resolution over  $A$  with augmentation  $\mu : P_0 \rightarrow A$ , and let  $i_0 : P_0 \rightarrow B_0(A)$  be the 0-th component of a morphism  $i_\bullet : P_\bullet \rightarrow B_\bullet(A)$  of complexes of  $A$ -bimodules lifting  $\mathrm{id}_A$ . The main aim of this subsection is

to prove the following theorem, that tells us that we can compute the Gerstenhaber bracket between  $\mathrm{HH}^0(A)$  and  $\mathrm{HH}^n(A)$  for  $n \in \mathbb{N}_0$  using a simple homological procedure on any projective bimodule resolution of  $A$ .

**THEOREM 2.7.** *Consider the same assumptions as in the previous paragraph. Let  $\eta_n : P_n \rightarrow P_n$  be the map given by  $\eta_n(v) = \rho v - v \rho$  for  $v \in P_n$  and  $n \in \mathbb{N}_0$ . Since  $\eta_\bullet = \{\eta_n : P_n \rightarrow P_n\}_{n \in \mathbb{N}_0}$  is a lifting of the zero morphism from  $A$  to itself,  $\eta_\bullet$  is null-homotopic, i.e. there is a family  $h_\bullet = \{h_n^\rho : P_n \rightarrow P_{n+1}\}_{n \in \mathbb{N}_0}$  of morphisms of  $A$ -bimodules such that*

$$\eta_0 = \partial_1 h_0^\rho \quad \text{and} \quad \eta_n = h_{n-1}^\rho \partial_n + \partial_{n+1} h_n^\rho \tag{2.3}$$

for  $n \in \mathbb{N}$ . Then, if  $\phi \in \mathrm{Hom}_{A^e}(P_n, A)$  is a cocycle for some  $n \in \mathbb{N}$ , the Gerstenhaber bracket  $[\phi, \ell_\rho i_0] \in \mathrm{HH}^{n-1}(A)$  is given by the cohomology class of  $\phi h_{n-1}^\rho$ .

**REMARK 2.8.** *It is easy to see that if  $\phi \in \mathrm{Hom}_{A^e}(P_n, A)$  is a cocycle (resp., coboundary), then  $\phi h_{n-1}^\rho$  is a cocycle (resp., coboundary) by applying (2.3). On the other hand, in general we have  $P_0 = B_0 = A \otimes A$  and  $i_0 = \mathrm{id}_{A \otimes A}$ , so we can forget about  $i_0$  in Theorem 2.7.*

The rest of this subsection is devoted to proving Theorem 2.7. In order to do that, we first need to prove some preliminary results.

Let  $t_n : B_n(A) \rightarrow B_{n+1}(A)$  be the morphism of  $A$ -bimodules given by

$$t_n(a_0 | \dots | a_{n+1}) = \sum_{j=0}^n (-1)^j a_0 | \dots | a_j | \rho | a_{j+1} | \dots | a_{n+1}$$

for  $a_0, \dots, a_{n+1} \in A$  and  $n \in \mathbb{N}_0$ . Let  $\xi_\bullet = \{\xi_n : B_n(A) \rightarrow B_n(A)\}_{n \in \mathbb{N}_0}$  be the family of morphisms of  $A$ -bimodules defined by  $\xi_n(u) = \rho u - u \rho$  for  $u \in B_n(A)$  and  $n \in \mathbb{N}_0$ .

**LEMMA 2.9.** *We have that  $\xi_0 = d_1 t_0$  and  $\xi_n = t_{n-1} d_n + d_{n+1} t_n$  for  $n \in \mathbb{N}$ .*

*Proof.* For  $a_0, \dots, a_{n+1} \in A$  and  $n \in \mathbb{N}$ ,

$$d_1 t_0(a_0 | a_1) = d_1(a_0 | \rho | a_1) = a_0 \rho | a_1 - a_0 | \rho a_1 = \rho a_0 | a_1 - a_0 | a_1 \rho = \xi_0(a_0 | a_1),$$

and

$$d_{n+1} t_n(a_0 | \dots | a_{n+1}) = d_{n+1} \left( \sum_{j=0}^n (-1)^j a_0 | \dots | a_j | \rho | a_{j+1} | \dots | a_{n+1} \right) = S_1 + S_2,$$

where

$$S_1 = \sum_{j=0}^n (-1)^j \left\{ (-1)^j a_0 | \dots | a_{j-1} | a_j \rho | a_{j+1} | \dots | a_{n+1} \right.$$

$$\begin{aligned} &+ (-1)^{j+1} a_0 | \dots | a_j | \rho a_{j+1} | a_{j+2} | \dots | a_{n+1} \Big\} \\ &= a_0 \rho | \dots | a_{n+1} - a_0 | \dots | a_n | \rho a_{n+1} \\ &= \xi_n(a_0 | \dots | a_{n+1}), \end{aligned}$$

and

$$\begin{aligned} S_2 &= \sum_{j=0}^n (-1)^j \left\{ \sum_{r=0}^{j-2} (-1)^r a_0 | \dots | a_{r-1} | a_r a_{r+1} | a_{r+2} | \dots | a_j | \rho | a_{j+1} | \dots | a_{n+1} \right. \\ &\quad + (-1)^{j-1} a_0 | \dots | a_{j-2} | a_{j-1} a_j | \rho | a_{j+1} | \dots | a_{n+1} \\ &\quad + (-1)^{j+2} a_0 | \dots | a_j | \rho | a_{j+1} a_{j+2} | a_{j+3} | \dots | a_{n+1} \\ &\quad \left. + \sum_{r=j+2}^n (-1)^{r+1} a_0 | \dots | a_j | \rho | a_{j+1} | \dots | a_{r-1} | a_r a_{r+1} | a_{r+2} | \dots | a_{n+1} \right\} \\ &= \sum_{i=0}^n \left\{ \sum_{j=i+2}^n (-1)^{j+i} a_0 | \dots | a_{i-1} | a_i a_{i+1} | a_{i+2} | \dots | a_j | \rho | a_{j+1} | \dots | a_{n+1} \right. \\ &\quad - a_0 | \dots | a_{i-1} | a_i a_{i+1} | \rho | a_{i+2} | \dots | a_{n+1} \\ &\quad + a_0 | \dots | a_{i-1} | \rho | a_i a_{i+1} | a_{i+2} | \dots | a_{n+1} \\ &\quad \left. + \sum_{j=0}^{i-2} (-1)^{j+i+1} a_0 | \dots | a_j | \rho | a_{j+1} | \dots | a_{i-1} | a_i a_{i+1} | a_{i+2} | \dots | a_{n+1} \right\} \\ &= \sum_{i=0}^n (-1)^{i+1} \left\{ \sum_{j=0}^{i-2} (-1)^j a_0 | \dots | a_j | \rho | a_{j+1} | \dots | a_{i-1} | a_i a_{i+1} | a_{i+2} | \dots | a_{n+1} \right. \\ &\quad + (-1)^{i-1} a_0 | \dots | a_{i-1} | \rho | a_i a_{i+1} | a_{i+2} | \dots | a_{n+1} \\ &\quad + (-1)^i a_0 | \dots | a_{i-1} | a_i a_{i+1} | \rho | a_{i+2} | \dots | a_{n+1} \\ &\quad \left. + \sum_{j=i+2}^n (-1)^{j-1} a_0 | \dots | a_{i-1} | a_i a_{i+1} | a_{i+2} | \dots | a_j | \rho | a_{j+1} | \dots | a_{n+1} \right\} \\ &= -t_{n-1} d_n(a_0 | \dots | a_{n+1}). \end{aligned}$$

Hence,  $\xi_n = t_{n-1} d_n + d_{n+1} t_n$ . □

LEMMA 2.10. *The Gerstenhaber bracket  $[\varphi, \ell_\rho] \in \mathrm{Hom}_{A^e}(B_{n-1}(A), A)$  is given by  $[\varphi, \ell_\rho] = \varphi t_{n-1}$  for  $\varphi \in \mathrm{Hom}_{A^e}(B_n(A), A)$  and  $n \in \mathbb{N}$ .*

*Proof.* For  $a_0, \dots, a_n \in A$ ,

$$\begin{aligned} [\varphi, \ell_\rho](a_0 | \dots | a_n) &= a_0 [F(\varphi), F(\ell_\rho)](a_1 | \dots | a_{n-1}) a_n \\ &= a_0 (F(\varphi) \circ_G F(\ell_\rho))(a_1 | \dots | a_{n-1}) a_n \\ &= a_0 \left( \sum_{i=1}^n (-1)^{i-1} F(\varphi)(a_1 | \dots | a_{i-1} | \rho | a_i | \dots | a_{n-1}) \right) a_n \end{aligned}$$

$$\begin{aligned}
 &= a_0 \left( \sum_{i=1}^n (-1)^{i-1} \varphi(1|a_1| \dots |a_{i-1}|\rho|a_i| \dots |a_{n-1}|1) \right) a_n \\
 &= \sum_{i=1}^n (-1)^{i-1} \varphi(a_0|a_1| \dots |a_{i-1}|\rho|a_i| \dots |a_{n-1}|a_n) \\
 &= (\varphi t_{n-1})(a_0|a_1| \dots |a_{n-1}|a_n).
 \end{aligned}$$

Hence,  $[\varphi, \ell_\rho] = \varphi t_{n-1}$ . □

LEMMA 2.11. *We assume the same hypotheses as those of Theorem 2.7. Then, there exists a family  $s_\bullet = \{s_n : P_n \rightarrow B_{n+2}(A)\}_{n \in \mathbb{N}_0}$  of morphisms of  $A$ -bimodules such that  $i_1 h_0^\rho - t_0 i_0 = d_2 s_0$  and  $i_{n+1} h_n^\rho - t_n i_n = d_{n+2} s_n - s_{n-1} \partial_n$  for  $n \in \mathbb{N}$ .*

*Proof.* Since  $d_1(i_1 h_0^\rho - t_0 i_0) = i_0 \partial_1 h_0^\rho - d_1 t_0 i_0 = i_0 \eta_0 - \xi_0 i_0 = 0$ , where we used that  $i_0$  is a morphism of  $A$ -bimodules in the last equality, there exists a morphism  $s_0 : P_0 \rightarrow B_2(A)$  of  $A$ -bimodules such that  $d_2 s_0 = i_1 h_0^\rho - t_0 i_0$ . We now claim that there exists a family  $s_\bullet = \{s_n : P_n \rightarrow B_{n+2}(A)\}_{n \in \mathbb{N}_0}$  of morphisms of  $A$ -bimodules such that  $d_{n+2} s_n = i_{n+1} h_n^\rho - t_n i_n + s_{n-1} \partial_n$  by induction on  $n \in \mathbb{N}_0$  (where  $s_{-1} = 0$ ). Indeed,

$$\begin{aligned}
 &d_{n+1}(i_{n+1} h_n^\rho - t_n i_n + s_{n-1} \partial_n) \\
 &= d_{n+1} i_{n+1} h_n^\rho - d_{n+1} t_n i_n + (i_n h_{n-1}^\rho - t_{n-1} i_{n-1} + s_{n-2} \partial_{n-1}) \partial_n \\
 &= i_n \partial_{n+1} h_n^\rho - (\xi_n - t_{n-1} d_n) i_n + i_n h_{n-1}^\rho \partial_n - t_{n-1} i_{n-1} \partial_n \\
 &= i_n (\partial_{n+1} h_n^\rho + h_{n-1}^\rho \partial_n) - \xi_n i_n = i_n \eta_n - \xi_n i_n = 0,
 \end{aligned}$$

where we used the inductive assumption in the first equality, Lemma 2.9 in the second equality, the definition of  $\eta_n$  in the third equality and the fact that  $i_n$  is a morphism of  $A$ -bimodules in the last equality. The result thus follows. □

*Proof of Theorem 2.7.* Let  $\varphi = \phi p_n \in \text{Hom}_{A^e}(B_n(A), A)$ . Then  $\varphi$  is a cocycle and  $[\phi, i_\bullet^*(\ell_\rho)] = i_\bullet^*([p_\bullet^*(\phi), \ell_\rho]) = [\varphi, \ell_\rho] i_{n-1} = \varphi t_{n-1} i_{n-1}$  by Lemma 2.10. Since  $p_\bullet i_\bullet$  is homotopic to the identity of  $P_\bullet$ , there exists  $\phi_1 \in \text{Hom}_{A^e}(P_{n-1}, A)$  such that  $\phi - \varphi i_n = \phi - \phi p_n i_n = \phi_1 \partial_n$ . Then,

$$\begin{aligned}
 \phi h_{n-1}^\rho - \varphi t_{n-1} i_{n-1} &= (\varphi i_n + \phi_1 \partial_n) h_{n-1}^\rho - \varphi t_{n-1} i_{n-1} \\
 &= \varphi (d_{n+1} s_{n-1} - s_{n-2} \partial_{n-1}) + \phi_1 (\eta_{n-1} - h_{n-2}^\rho \partial_{n-1}) \\
 &= -\varphi s_{n-2} \partial_{n-1} - \phi_1 h_{n-2}^\rho \partial_{n-1} \in \text{Hom}_{A^e}(P_{n-1}, A)
 \end{aligned}$$

is a boundary, where we used Lemma 2.11 and the definition of  $\eta_{n-1}$  in the second equality and the fact that  $\phi_1$  is a morphism of  $A$ -bimodules in the last identity. Hence,  $[\phi, i_\bullet^*(\ell_\rho)] \in \text{HH}^{n-1}(A)$  coincides with the cohomology class of  $\phi h_{n-1}^\rho$ , as was to be shown. □

REMARK 2.12. *The homotopy maps  $h_\bullet^\rho$  in Theorem 2.7 are presumably homotopy liftings in the sense of [9]. However, our maps  $h_\bullet^\rho$  do not directly follow the*



scheme of that definition –as well as being far simpler, for they are restricted to a much easier situation– since they do not require the computation of any map  $\Delta : P_\bullet \rightarrow P_\bullet \otimes_A P_\bullet$  lifting the isomorphism  $A \rightarrow A \otimes_A A$ , which is also the case in [6].

2.3 METHOD COMPUTING THE BRACKET BETWEEN  $\mathrm{HH}^1(A)$  AND  $\mathrm{HH}^n(A)$   
(AFTER M. SUÁREZ-ÁLVAREZ)

In this subsection we will briefly recall the method introduced by M. Suárez-Álvarez in [8] to compute the Gerstenhaber bracket between  $\mathrm{HH}^1(A)$  and  $\mathrm{HH}^n(A)$  for  $n \in \mathbb{N}_0$ .

Recall that  $\mathrm{HH}^1(A)$  is isomorphic to the quotient of the space of derivations of  $A$  modulo the subspace of inner derivations. Let  $\rho : A \rightarrow A$  be a derivation of  $A$ , i.e.  $\rho(xy) = \rho(x)y + x\rho(y)$  for all  $x, y \in A$ . For a left  $A$ -module  $M$ , a  $\rho$ -OPERATOR on  $M$  is a map  $f : M \rightarrow M$  such that  $f(am) = \rho(a)m + af(m)$  for all  $a \in A$  and  $m \in M$ . It is direct to see that the map  $\rho^e = \rho \otimes \mathrm{id}_A + \mathrm{id}_A \otimes \rho : A^e \rightarrow A^e$  defined by  $\rho^e(x \otimes y) = \rho(x) \otimes y + x \otimes \rho(y)$  for  $x, y \in A$  is a derivation of the enveloping algebra  $A^e$  and  $\rho$  is a  $\rho^e$ -operator on  $A$ .

Let  $(P_\bullet, \partial_\bullet)$  be a projective bimodule resolution over  $A$  with augmentation  $\mu : P_0 \rightarrow A$ . A  $\rho^e$ -LIFTING of  $\rho$  to  $(P_\bullet, \partial_\bullet)$  is a family of  $\rho^e$ -operators  $\rho_\bullet = \{\rho_n : P_n \rightarrow P_n\}_{n \in \mathbb{N}_0}$  such that  $\mu\rho_0 = \rho\mu$  and  $\partial_n\rho_n = \rho_{n-1}\partial_n$  for  $n \in \mathbb{N}$ . The morphism of complexes

$$\rho_{\bullet, P_\bullet}^\sharp : \mathrm{Hom}_{A^e}(P_\bullet, A) \rightarrow \mathrm{Hom}_{A^e}(P_\bullet, A)$$

defined by  $\rho_{n, P_\bullet}^\sharp(\phi) = \rho\phi - \phi\rho_n$  for  $\phi \in \mathrm{Hom}_{A^e}(P_n, A)$  and  $n \in \mathbb{N}_0$  is independent of the  $\rho^e$ -lifting up to homotopy (see [8], Lemma 1.6) and it thus induces a morphism on cohomology that we will denote by the same symbol. Let  $i_\bullet : P_\bullet \rightarrow B_\bullet(A)$  and  $p_\bullet : B_\bullet(A) \rightarrow P_\bullet$  be morphisms of complexes of  $A$ -bimodules lifting  $\mathrm{id}_A$ . Then the diagram

$$\begin{CD} \mathrm{H}^n(\mathrm{Hom}_{A^e}(B_\bullet(A), A)) @>{\mathrm{H}(\rho_{\bullet, B_\bullet(A)}^\sharp)}>> \mathrm{H}^n(\mathrm{Hom}_{A^e}(B_\bullet(A), A)) \\ @VV{\mathrm{H}(i_\bullet^\sharp)}V @VV{\mathrm{H}(i_\bullet^\sharp)}V \\ \mathrm{H}^n(\mathrm{Hom}_{A^e}(P_\bullet, A)) @>{\mathrm{H}(\rho_{\bullet, P_\bullet}^\sharp)}>> \mathrm{H}^n(\mathrm{Hom}_{A^e}(P_\bullet, A)) \end{CD} \tag{2.4}$$

commutes (see [8], Lemma 1.6). On the other hand, as noted in [8], Sections 2.1 and 2.2, using the  $\rho^e$ -lifting of  $\rho$  to the bar resolution defined by

$$\rho_n(a_0 | \dots | a_{n+1}) = \sum_{j=0}^{n+1} a_0 | \dots | a_{j-1} | \rho(a_j) | a_{j+1} | \dots | a_{n+1}$$

for  $a_0, \dots, a_{n+1} \in A$  and  $n \in \mathbb{N}_0$ , it is easy to check that the diagram

$$\begin{CD} \mathrm{Hom}_{A^e}(B_n(A), A) @>\rho_{n, B_\bullet(A)}^\sharp>> \mathrm{Hom}_{A^e}(B_n(A), A) \\ @V F VV @VV F V \\ \mathrm{Hom}_{\mathbb{k}}(A^{\otimes n}, A) @>[\rho, -]>> \mathrm{Hom}_{\mathbb{k}}(A^{\otimes n}, A) \end{CD}$$

commutes. As a consequence, the Gerstenhaber bracket between the cohomology classes of  $G(\rho) \in \mathrm{Hom}_{A^e}(B_1(A), A)$  and  $\varphi \in \mathrm{Hom}_{A^e}(B_n(A), A)$  is given by the cohomology class of  $[G(\rho), \varphi] = G([\rho, F(\varphi)]) = \rho_{n, B_\bullet(A)}^\sharp(\varphi)$ .

We finally recall one of the main results of [8], which tells us that we can compute the Gerstenhaber bracket between  $\mathrm{HH}^1(A)$  and  $\mathrm{HH}^n(A)$  for  $n \in \mathbb{N}_0$  using any projective bimodule resolution of  $A$  (see [8], Thm. A and Section 2.2). The proof just follows from observing that, on cohomology, (2.4) gives us the identities

$$[i_\bullet^*(G(\rho)), \phi] = i_\bullet^*([G(\rho), p_\bullet^*(\phi)]) = i_\bullet^*(\rho_{n, B_\bullet(A)}^\sharp(p_\bullet^*(\phi))) = \rho_{n, P_\bullet}^\sharp(\phi).$$

**THEOREM 2.13.** *Let  $(P_\bullet, \partial_\bullet)$  be a projective bimodule resolution over the algebra  $A$  with augmentation  $\mu : P_0 \rightarrow A$ , and let  $i_1 : P_1 \rightarrow B_1(A)$  be the first component of the morphism  $i_\bullet : P_\bullet \rightarrow B_\bullet(A)$  of complexes of  $A$ -bimodules lifting  $\mathrm{id}_A$ . Given a cocycle  $\phi \in \mathrm{Hom}_{A^e}(P_n, A)$  and  $n \in \mathbb{N}_0$ , the Gerstenhaber bracket  $[G(\rho)i_1, \phi] \in \mathrm{HH}^n(A)$  is given by the cohomology class of  $\rho_{n, P_\bullet}^\sharp(\phi)$ .*

**REMARK 2.14.** *Note that in our Theorem 2.7, as well as in the result proved in [8] that was recalled before as Theorem 2.13, we need at least some component(s) of the comparison map from the generic projective resolution  $(P_\bullet, \partial_\bullet)$  to the bar resolution.*

### 3 BASICS ON THE FOMIN-KIRILLOV ALGEBRA $\mathrm{FK}(3)$ ON 3 GENERATORS

In this section we will review the basic definitions concerning the Fomin-Kirillov algebra on 3 generators. Recall that, given  $i \in \mathbb{Z}$ , we will denote by  $\mathbb{Z}_{\leq i}$  the set  $\{m \in \mathbb{Z} | m \leq i\}$ . Given  $r \in \mathbb{R}$ , we set  $\lfloor r \rfloor = \sup\{n \in \mathbb{Z} | n \leq r\}$  the usual FLOOR FUNCTION. From now on,  $\mathbb{k}$  is a field of characteristic different from 2 and 3. We recall that the FOMIN-KIRILLOV ALGEBRA on 3 generators is the  $\mathbb{k}$ -algebra  $\mathrm{FK}(3)$  generated by the  $\mathbb{k}$ -vector space  $V$  spanned by three elements  $a, b$  and  $c$  modulo the ideal generated by the vector space  $R \subseteq V^{\otimes 2}$  spanned by

$$\{a^2, b^2, c^2, ab + bc + ca, ba + ac + cb\}.$$

This is a nonnegatively graded algebra by setting the generators  $a, b$  and  $c$  in (internal) degree 1. As usual, we will omit the tensor symbol  $\otimes$  when denoting the product of the elements of the tensor algebra  $\mathrm{TV} = \bigoplus_{n \in \mathbb{N}_0} V^{\otimes n}$ . We refer the reader to [1, 4] for more information on Fomin-Kirillov algebras.

Note that  $\mathrm{FK}(3) = \bigoplus_{m \in \llbracket 0, 4 \rrbracket} \mathrm{FK}(3)_m$ , where  $\mathrm{FK}(3)_m$  is the subspace of  $\mathrm{FK}(3)$  concentrated in internal degree  $m$ . It is easy to see that

$$\mathcal{B} = \{1, a, b, c, ab, bc, ba, ac, aba, abc, bac, abac\} \tag{3.1}$$

is a basis of  $\mathrm{FK}(3)$  (see [1]). Given  $m \in \llbracket 0, 4 \rrbracket$ , we will denote by  $\mathcal{B}_m$  the subset of (3.1) that is a basis of  $\mathrm{FK}(3)_m$ .

Let us briefly denote by  $\{A, B, C\}$  the basis of  $V^*$  dual to the basis  $\{a, b, c\}$  of  $V$ , where the former are concentrated in internal degree  $-1$ . The **QUADRATIC DUAL**  $\mathrm{FK}(3)^! = \bigoplus_{n \in \mathbb{N}_0} \mathrm{FK}(3)^!_{-n}$  of the Fomin Kirillov algebra  $\mathrm{FK}(3)$  is then given by

$$\mathrm{FK}(3)^! = \mathbb{k}\langle A, B, C \rangle / (BA - AC, CA - AB, AB - BC, CB - BA),$$

where  $\mathrm{FK}(3)^!_{-n}$  is the subspace of  $\mathrm{FK}(3)^!$  concentrated in internal degree  $-n$ . Notice that  $\mathrm{FK}(3)^!_0 = \mathbb{k}$  and  $\mathrm{FK}(3)^!_{-1} \cong V^*$ . We recall that  $\mathcal{B}_n^! = \{A^n, B^n, C^n, A^{n-1}B, A^{n-1}C, A^{n-2}B^2\}$  is a basis of  $\mathrm{FK}(3)^!_{-n}$  for all integers  $n \geq 2$ , where we follow the convention that  $A^0B^2 = B^2$  (see [7], Lemma 4.4).

FOR SIMPLICITY, FROM NOW ON WE WILL DENOTE THE FOMIN-KIRILLOV ALGEBRA  $\mathrm{FK}(3)$  SIMPLY BY  $A$ . Let  $(A^!_{-n})^*$  be the dual space of  $A^!_{-n}$  and  $\mathcal{B}_n^!^* = \{\alpha_n, \beta_n, \gamma_n, \alpha_{n-1}\beta, \alpha_{n-1}\gamma, \alpha_{n-2}\beta^2\} \setminus \{\mathbf{0}\}$  the dual basis to  $\mathcal{B}_n^!$  for  $n \in \mathbb{N}$ , where we will follow the convention that if the index of some letter in an element of the previous sets is less than or equal to zero, it is the zero element  $\mathbf{0}$ . We will omit the index 1 for the elements of the previous bases and write  $\mathcal{B}_0^!^* = \{e^!\}$ , where  $e^!$  is the basis element of  $(A^!_0)^*$ . The previous bases for the homogeneous components of  $A$  or  $(A^!)^\# = \bigoplus_{n \in \mathbb{N}_0} (A^!_{-n})^*$  will be called **USUAL**. Recall that  $(A^!)^\#$  is a graded bimodule over  $A^!$  via  $(ufv)(w) = f(vwu)$  for  $u, v, w \in A^!$  and  $f \in (A^!)^\#$ .

For the explicit description of the **BIMODULE KOSZUL COMPLEX**  $(K_\bullet^b, d_\bullet^b)$  over the Fomin-Kirillov algebra  $A$ , we refer the reader to [3], Subsection 3.1. The following result gives an explicit description of the minimal projective resolution of  $A$  in the category of bounded below graded  $A$ -bimodules.

**PROPOSITION 3.1.** ([3], Prop. 3.7) *The minimal projective resolution  $(P_\bullet^b, \delta_\bullet^b)$  of  $A$  in the category of bounded below graded  $A$ -bimodules is given as follows. For  $n \in \mathbb{N}_0$ , set*

$$P_n^b = \bigoplus_{i \in \llbracket 0, \lfloor n/4 \rfloor \rrbracket} \omega_i K_{n-4i}^b = \bigoplus_{i \in \llbracket 0, \lfloor n/4 \rfloor \rrbracket} \omega_i A \otimes (A^!_{-(n-4i)})^* \otimes A,$$

where  $\omega_i$  is a symbol of homological degree  $4i$  and internal degree  $6i$  for all  $i \in \mathbb{N}_0$ , the  $A$ -bimodule structure of  $P_n^b$  is given by  $x'(\omega_i x \otimes u \otimes y)y' = \omega_i x'x \otimes u \otimes yy'$  for all  $x, x', y, y' \in A$  and  $u \in (A^!_{-(n-4i)})^*$ , and the differential  $\delta_n^b : P_n^b \rightarrow P_{n-1}^b$  for  $n \in \mathbb{N}$  is given by

$$\delta_n^b \left( \sum_{i \in \llbracket 0, \lfloor n/4 \rfloor \rrbracket} \omega_i \rho_{n-4i} \right) = \sum_{i \in \llbracket 0, \lfloor n/4 \rfloor \rrbracket} (\omega_i d_{n-4i}^b(\rho_{n-4i}) + \omega_{i-1} f_{n-4i}^b(\rho_{n-4i})),$$

where  $\rho_j \in K_j^b$  for  $j \in \mathbb{N}_0$ ,  $\omega_{-1} = 0$  and  $f_j^b : K_j^b \rightarrow K_{j+3}^b$  are the morphisms given in [3], (3.2). This gives a minimal projective resolution of  $A$  by means of the augmentation  $\epsilon^b : P_0^b = A \otimes (A_0^!)^* \otimes A \rightarrow A$ , where  $\epsilon^b(x \otimes \epsilon^! \otimes y) = xy$  for  $x, y \in A$ .

To reduce space, we will typically use vertical bars instead of the tensor product symbols  $\otimes$  for the elements of  $P_\bullet$ . We will use in the sequel the following explicit expression of some values of the map  $f_0^b$  in the previous result. They follow immediately from [3], (3.2).

FACT 3.2. *The morphism  $f_0^b$  defined in [3], (3.2), satisfies that*

$$\begin{aligned}
 f_0^b(a|\epsilon^!|1) &= 2a|\alpha_3|bac + 2a|\beta_3|abc - 2a|\gamma_3|aba - a|\alpha_2\beta|abc + a|\alpha_2\gamma|aba \\
 &\quad - a|\alpha\beta_2|bac - ac|\alpha_2\beta|ab - ab|\alpha_2\gamma|ac + ab|\alpha\beta_2|ab + ab|\alpha\beta_2|bc \\
 &\quad - ac|\alpha\beta_2|ba - 2ab|\alpha_3|ab - 2ab|\alpha_3|bc + 2ac|\alpha_3|ba + 2ac|\beta_3|ab \\
 &\quad + 2ab|\gamma_3|ac - abc|\alpha_2\beta|a - aba|\alpha_2\beta|c + abc|\alpha_2\gamma|b + aba|\alpha_2\gamma|a \\
 &\quad + 2abc|\beta_3|a + 2aba|\beta_3|c - 2abc|\gamma_3|b - 2aba|\gamma_3|a + 2abac|\alpha_3|1 \\
 &\quad - abac|\alpha\beta_2|1, \\
 f_0^b(1|\epsilon^!|a) &= -2|\alpha_3|abac + 1|\alpha\beta_2|abac - a|\alpha_2\beta|abc - c|\alpha_2\beta|aba + a|\alpha_2\gamma|aba \\
 &\quad + b|\alpha_2\gamma|abc + 2a|\beta_3|abc + 2c|\beta_3|aba - 2a|\gamma_3|aba - 2b|\gamma_3|abc \\
 &\quad + ba|\alpha_2\beta|(ab + bc) + ab|\alpha_2\gamma|ba + bc|\alpha_2\gamma|ba + ab|\alpha\beta_2|(ab + bc) \\
 &\quad - ac|\alpha\beta_2|ba - 2ab|\alpha_3|(ab + bc) + 2ac|\alpha_3|ba - 2ba|\beta_3|(ab + bc) \\
 &\quad - 2ab|\gamma_3|ba - 2bc|\gamma_3|ba + 2bac|\alpha_3|a + 2abc|\beta_3|a - 2aba|\gamma_3|a \\
 &\quad - abc|\alpha_2\beta|a + aba|\alpha_2\gamma|a - bac|\alpha\beta_2|a, \\
 f_0^b(b|\epsilon^!|1) &= 2b|\alpha_3|bac + 2b|\beta_3|abc - 2b|\gamma_3|aba - b|\alpha_2\beta|abc + b|\alpha_2\gamma|aba \\
 &\quad - b|\alpha\beta_2|bac + ba|\alpha_2\beta|ac + ba|\alpha_2\beta|ba - bc|\alpha_2\beta|ab - ba|\alpha_2\gamma|bc \\
 &\quad - bc|\alpha\beta_2|ba + 2bc|\alpha_3|ba - 2ba|\beta_3|ac - 2ba|\beta_3|ba + 2bc|\beta_3|ab \\
 &\quad + 2ba|\gamma_3|bc + aba|\alpha_2\gamma|b + bac|\alpha_2\gamma|a - aba|\alpha\beta_2|c - bac|\alpha\beta_2|b \\
 &\quad + 2aba|\alpha_3|c + 2bac|\alpha_3|b - 2aba|\gamma_3|b - 2bac|\gamma_3|a + 2abac|\beta_3|1 \\
 &\quad - abac|\alpha_2\beta|1, \\
 f_0^b(1|\epsilon^!|b) &= -2|\beta_3|abac + 1|\alpha_2\beta|abac + a|\alpha_2\gamma|bac + b|\alpha_2\gamma|aba - b|\alpha\beta_2|bac \\
 &\quad - c|\alpha\beta_2|aba + 2b|\alpha_3|bac + 2c|\alpha_3|aba - 2a|\gamma_3|bac - 2b|\gamma_3|aba \\
 &\quad - bc|\alpha_2\beta|ab + ba|\alpha_2\beta|(ba + ac) + ba|\alpha_2\gamma|ab + ac|\alpha_2\gamma|ab \\
 &\quad + ab|\alpha\beta_2|(ba + ac) - 2ab|\alpha_3|(ba + ac) + 2bc|\beta_3|ab \\
 &\quad - 2ba|\beta_3|(ba + ac) - 2ba|\gamma_3|ab - 2ac|\gamma_3|ab + 2bac|\alpha_3|b + 2abc|\beta_3|b \\
 &\quad - 2aba|\gamma_3|b - abc|\alpha_2\beta|b + aba|\alpha_2\gamma|b - bac|\alpha\beta_2|b, \\
 f_0^b(c|\epsilon^!|1) &= 2c|\alpha_3|bac + 2c|\beta_3|abc - 2c|\gamma_3|aba - c|\alpha_2\beta|abc + c|\alpha_2\gamma|aba \\
 &\quad - c|\alpha\beta_2|bac - (ab + bc)|\alpha_2\beta|ac - (ab + bc)|\alpha_2\beta|ba \\
 &\quad + (ab + bc)|\alpha_2\gamma|bc + (ba + ac)|\alpha_2\gamma|ac - (ba + ac)|\alpha\beta_2|ab
 \end{aligned}$$

$$\begin{aligned}
 & - (ba + ac)|\alpha\beta_2|bc + 2(ba + ac)|\alpha_3|ab + 2(ba + ac)|\alpha_3|bc \\
 & + 2(ab + bc)|\beta_3|ac + 2(ab + bc)|\beta_3|ba - 2(ab + bc)|\gamma_3|bc \\
 & - 2(ba + ac)|\gamma_3|ac + bac|\alpha_2\beta|a - abc|\alpha_2\beta|c - bac|\alpha\beta_2|c \\
 & + abc|\alpha\beta_2|b + 2bac|\alpha_3|c - 2abc|\alpha_3|b - 2bac|\beta_3|a + 2abc|\beta_3|c \\
 & + 2abac|\gamma_3|1 - abac|\alpha_2\gamma|1, \\
 f_0^b(1|\epsilon^1|c) = & -2|\gamma_3|abac + 1|\alpha_2\gamma|abac + a|\alpha_2\beta|bac - c|\alpha_2\beta|abc + b|\alpha\beta_2|abc \\
 & - c|\alpha\beta_2|bac - 2b|\alpha_3|abc + 2c|\alpha_3|bac - 2a|\beta_3|bac + 2c|\beta_3|abc \\
 & - bc|\alpha_2\beta|ac + ab|\alpha_2\gamma|bc + bc|\alpha_2\gamma|bc + ba|\alpha_2\gamma|ac + ac|\alpha_2\gamma|ac \\
 & - ac|\alpha\beta_2|bc + 2ac|\alpha_3|bc + 2bc|\beta_3|ac - 2ab|\gamma_3|bc - 2bc|\gamma_3|bc \\
 & - 2ba|\gamma_3|ac - 2ac|\gamma_3|ac + 2bac|\alpha_3|c + 2abc|\beta_3|c - 2aba|\gamma_3|c \\
 & - abc|\alpha_2\beta|c + aba|\alpha_2\gamma|c - bac|\alpha\beta_2|c.
 \end{aligned}$$

Given  $n \in \mathbb{N}_0$ , there is a canonical isomorphism

$$\mathrm{Hom}_{A^e}(P_n^b, A) \cong Q^n \tag{3.2}$$

of graded  $\mathbb{k}$ -vector spaces, where  $Q^n = \bigoplus_{i \in [0, \lfloor n/4 \rfloor]} \omega_i^* K^{n-4i}$  and  $K^n = \mathrm{Hom}_{\mathbb{k}}((A_{-n}^1)^*, A)$ . Transporting the differential of the left member of (3.2) induced by that of  $(P_\bullet^b, \delta_\bullet^b)$ , we obtain a complex of  $\mathbb{k}$ -vector spaces  $Q^\bullet$ , whose cohomology gives the linear structure of the Hochschild cohomology  $\mathrm{HH}^\bullet(A)$ . Note that the space  $\mathrm{Hom}_{\mathbb{k}}((A_{-n}^1)^*, A_m)$  is concentrated in cohomological degree  $n$  and internal degree  $m - n$ . The symbol  $\omega_i^*$  has cohomological degree  $4i$  and internal degree  $-6i$  for  $i \in \mathbb{N}_0$ . We will usually omit  $\omega_0^*$  for simplicity. Let  $H_m^n$  be the subspace of  $H^n(Q^\bullet)$  concentrated in internal degree  $m - n$  for  $m, n \in \mathbb{Z}$ . Note that  $H_m^n = 0$  for  $(n, m) \in \mathbb{Z}^2 \setminus (\mathbb{N}_0 \times \mathbb{Z}_{\leq 4})$ . The following result gives a recursive description of the spaces  $H_m^n$ .

PROPOSITION 3.3. ([3], Cor. 5.3) *For integers  $m \leq 1$  and  $n \in \mathbb{N}_0$ , we have*

$$H_m^n \cong \begin{cases} \omega_{\frac{1-m}{2}}^* H_1^{n+2m-2}, & \text{if } m \text{ is odd,} \\ \omega_{-\frac{m}{2}}^* H_0^{n+2m}, & \text{if } m \text{ is even.} \end{cases}$$

Given elements  $x \in \mathfrak{B}_m$  and  $y \in \mathfrak{B}_n^*$ , the symbol  $y|x$  will denote the linear map in  $K^n = \mathrm{Hom}_{\mathbb{k}}((A_{-n}^1)^*, A)$ , which maps  $y$  to  $x$  and sends the other usual basis elements of  $(A_{-n}^1)^*$  to zero. See [3], Cor. 5.4, for specific representatives of the cohomology classes of a basis of  $H_m^n$  for  $(n, m) \in \mathbb{N}_0 \times \mathbb{Z}_{\leq 4}$ .

The algebra structure of the Hochschild cohomology  $\mathrm{HH}^\bullet(A)$  (with the multiplication given by the cup product) is described as follows.

THEOREM 3.4. ([3], Cor. 6.11) *The Hochschild cohomology  $\mathrm{HH}^\bullet(A)$  is isomorphic to the quotient of the free graded-commutative (for the cohomological degree)  $\mathbb{k}$ -algebra generated by fourteen elements (with fixed cohomological degrees and internal degrees) modulo the ideal generated by the elements given in [3], (6.5).*

Moreover, the fourteen generators of  $\mathrm{HH}^\bullet(A)$  mentioned in Theorem 3.4 are represented in  $\mathrm{H}^\bullet(Q^\bullet)$  by the following cocycles:  $X_1 = \epsilon^1(ab + ba)$ ,  $X_2 = \epsilon^1(ab + bc - ac)$ ,  $X_3 = \epsilon^1|abac$ ,  $X_4 = \alpha|bac$ ,  $X_5 = \beta|abc$ ,  $X_6 = \gamma|aba$ ,  $X_7 = \alpha|(aba - abc)$ ,  $X_8 = \alpha|a + \beta|b + \gamma|c$ ,  $X_9 = \alpha_2|1$ ,  $X_{10} = \beta_2|1$ ,  $X_{11} = \gamma_2|1$ ,  $X_{12} = (\alpha\beta + \alpha\gamma)|1$ ,  $X_{13} = \alpha_3|a + \beta_3|b + \gamma_3|c$  and  $X_{14} = \omega_1^*\epsilon^1|1$ . Let  $Y_i \in \mathrm{Hom}_{A^e}(P_n^b, A)$  be the element associated to  $X_i$  via the isomorphism (3.2) for  $i \in \llbracket 1, 14 \rrbracket$ . In what follows and to simplify our notation, given a cocycle  $\phi$ , we will use the same symbol  $\phi$  for its cohomology class.

Let  $i_\bullet : P_\bullet^b \rightarrow B_\bullet(A)$  be a morphism of complexes of  $A$ -bimodules lifting  $\mathrm{id}_A$ . It is clear that  $i_0 : A \otimes (A_0^1)^* \otimes A \rightarrow A \otimes A$  and  $i_1 : A \otimes (A_{-1}^1)^* \otimes A \rightarrow A^{\otimes 3}$  can be chosen as follows

$$i_0(1|\epsilon^1|1) = 1|1, i_1(1|\alpha|1) = -1|a|1, i_1(1|\beta|1) = -1|b|1 \text{ and } i_1(1|\gamma|1) = -1|c|1.$$

#### 4 GERSTENHABER BRACKETS ON HOCHSCHILD COHOMOLOGY OF $\mathrm{FK}(3)$

##### 4.1 GERSTENHABER BRACKETS OF $\mathrm{HH}^0(A)$ WITH $\mathrm{HH}^n(A)$

In this subsection we are going to utilize the method introduced in Subsection 2.2 to compute the Gerstenhaber bracket of  $X_i$  for  $i \in \llbracket 1, 14 \rrbracket$  with the elements  $X_1, X_2, X_3$  in  $\mathrm{HH}^0(A)$ . To wit, for every element  $X_i$  with  $i \in \llbracket 1, 3 \rrbracket$ , we find the associated element  $\rho$  in the center  $\mathcal{Z}(A)$  such that  $\ell_\rho i_0 = X_i$ , provide the corresponding self-homotopy  $h_\bullet^\rho$  satisfying (2.3) and then compute the respective Gerstenhaber brackets by means of Theorem 2.7.

We remark first that  $[X_i, 1] = 0$  for  $i \in \llbracket 1, 14 \rrbracket$ , since  $h_\bullet^1 = 0$  gives  $[X_i, 1] = 0$  for  $i \in \llbracket 4, 14 \rrbracket$  and the other follow from cohomological degree reasons. On the other hand, Definition 2.1 tells us that  $[X_i, X_j] = 0$  for  $i, j \in \llbracket 1, 3 \rrbracket$ . The proof of the following three results is a lengthy but straightforward computation.

**FACT 4.1.** *Let  $\rho = ab + ba \in \mathcal{Z}(A)$ . Then, there is a self-contracting homotopy  $h_\bullet^\rho$  satisfying (2.3) such that*

$$\begin{aligned} h_0^\rho(1|\epsilon^1|1) &= -b|\alpha|1 - a|\beta|1 - 1|\alpha|b - 1|\beta|a, \\ h_n^\rho(1|\alpha_n|1) &= (-1)^{n+1}b|\alpha_{n+1}|1 - 1|\alpha_{n+1}|b, \\ h_n^\rho(1|\beta_n|1) &= (-1)^{n+1}a|\beta_{n+1}|1 - 1|\beta_{n+1}|a \end{aligned}$$

for  $n \in \mathbb{N}$ , and

$$\begin{aligned} h_1^\rho(1|\gamma|1) &= b|\alpha_2|1 + a|\beta_2|1 + a|\alpha\beta|1 + b|\alpha\gamma|1 - 1|\alpha_2|b - 1|\beta_2|a - 1|\alpha\beta|b \\ &\quad - 1|\alpha\gamma|a, \\ h_2^\rho(1|\gamma_2|1) &= a|\gamma_3|1 + b|\gamma_3|1 + c|\alpha_2\beta|1 + c|\alpha\beta_2|1 + 1|\gamma_3|a + 1|\gamma_3|b + 1|\alpha_2\beta|c \\ &\quad + 1|\alpha\beta_2|c, \\ h_2^\rho(1|\alpha\beta|1) &= -b|\alpha_3|1 - c|\beta_3|1 - a|\alpha_2\gamma|1 - 1|\alpha_3|c - 1|\beta_3|a - 1|\alpha_2\gamma|b, \\ h_2^\rho(1|\alpha\gamma|1) &= -c|\alpha_3|1 - a|\beta_3|1 - b|\alpha_2\gamma|1 - 1|\alpha_3|b - 1|\beta_3|c - 1|\alpha_2\gamma|a. \end{aligned}$$

FACT 4.2. *Let  $\rho = ab + bc - ac \in \mathcal{Z}(A)$ . Then, there is a self-contracting homotopy  $h_\bullet^\rho$  satisfying (2.3) such that*

$$\begin{aligned} h_0^\rho(1|\epsilon^1|1) &= c|\alpha|1 + a|\gamma|1 + 1|\alpha|c + 1|\gamma|a, \\ h_n^\rho(1|\alpha_n|1) &= (-1)^n c|\alpha_{n+1}|1 + 1|\alpha_{n+1}|c, \\ h_n^\rho(1|\gamma_n|1) &= (-1)^n a|\gamma_{n+1}|1 + 1|\gamma_{n+1}|a \end{aligned}$$

for  $n \in \mathbb{N}$ , and

$$\begin{aligned} h_1^\rho(1|\beta|1) &= -c|\alpha_2|1 - a|\gamma_2|1 - c|\alpha\beta|1 - a|\alpha\gamma|1 + 1|\alpha_2|c + 1|\gamma_2|a + 1|\alpha\beta|a \\ &\quad + 1|\alpha\gamma|c, \\ h_2^\rho(1|\beta_2|1) &= -a|\beta_3|1 - c|\beta_3|1 - b|\alpha_2\gamma|1 - b|\alpha\beta_2|1 - 1|\beta_3|a - 1|\beta_3|c - 1|\alpha_2\gamma|b \\ &\quad - 1|\alpha\beta_2|b, \\ h_2^\rho(1|\alpha\beta|1) &= b|\alpha_3|1 + a|\gamma_3|1 + c|\alpha_2\beta|1 + 1|\alpha_3|c + 1|\gamma_3|b + 1|\alpha_2\beta|a, \\ h_2^\rho(1|\alpha\gamma|1) &= c|\alpha_3|1 + b|\gamma_3|1 + a|\alpha_2\beta|1 + 1|\alpha_3|b + 1|\gamma_3|a + 1|\alpha_2\beta|c. \end{aligned}$$

FACT 4.3. *Let  $\rho = abac \in \mathcal{Z}(A)$ . Then, there is a self-contracting homotopy  $h_\bullet^\rho$  satisfying (2.3) such that*

$$\begin{aligned} h_0^\rho(1|\epsilon^1|1) &= -aba|\gamma|1 - ab|\alpha|c - a|\beta|ac - 1|\alpha|bac, \\ h_1^\rho(1|\alpha|1) &= aba|\alpha\beta|1 - ab|\alpha_2|b - ba|\beta_2|c + c|\alpha_2|bc + b|\beta_2|ac + b|\alpha\beta|bc \\ &\quad - 1|\alpha_2|bac - 1|\alpha\beta|abc, \\ h_1^\rho(1|\beta|1) &= aba|\alpha\gamma|1 - 2ab|\alpha_2|c - ac|\alpha_2|a - ab|\alpha\beta|a + a|\alpha_2|bc - a|\beta_2|ab \\ &\quad - a|\beta_2|bc + c|\beta_2|ac + a|\alpha\gamma|ac - 1|\alpha\gamma|bac, \\ h_1^\rho(1|\gamma|1) &= 2aba|\gamma_2|1 - ba|\alpha\beta|c + b|\alpha_2|bc - a|\gamma_2|ac - c|\alpha\beta|ac - 1|\alpha\gamma|abc, \\ h_2^\rho(1|\alpha_2|1) &= bac|\alpha_3|1 + bc|\beta_3|a - ba|\beta_3|c + ba|\alpha_2\gamma|a - b|\alpha_3|ab - b|\alpha_3|bc \\ &\quad + c|\alpha_3|ba + a|\beta_3|ac + c|\beta_3|bc + a|\alpha_2\gamma|bc + b|\alpha_2\gamma|ba - 2|\alpha_3|bac, \\ h_2^\rho(1|\beta_2|1) &= abc|\beta_3|1 - 2ab|\alpha_3|c + ac|\alpha_3|b + ab|\beta_3|a - bc|\beta_3|a + ab|\alpha_2\gamma|b \\ &\quad - ba|\alpha_2\gamma|a + b|\alpha_3|ab + 2b|\alpha_3|bc - c|\alpha_3|ba + c|\alpha_3|ac - a|\beta_3|ac \\ &\quad - b|\alpha_2\gamma|ba + b|\alpha_2\gamma|ac - 1|\alpha_3|bac - 2|\beta_3|abc - 1|\alpha_2\gamma|aba, \\ h_2^\rho(1|\gamma_2|1) &= -3aba|\gamma_3|1 + ba|\beta_3|c - ab|\alpha\beta_2|c - b|\alpha_3|bc - a|\beta_3|ac - c|\beta_3|bc \\ &\quad + b|\gamma_3|ac - a|\alpha_2\gamma|bc + 1|\alpha_3|bac - 1|\alpha_2\beta|abc, \\ h_2^\rho(1|\alpha\beta|1) &= -2aba|\alpha\beta_2|1 - ac|\alpha_3|c - bc|\beta_3|b - 2ba|\beta_3|a - ab|\alpha_2\gamma|c \\ &\quad - ba|\alpha_2\gamma|b - a|\alpha_3|bc + b|\alpha_3|ba + 2b|\alpha_3|ac + b|\beta_3|ab - c|\beta_3|ac \\ &\quad + a|\gamma_3|ac - c|\gamma_3|ab + c|\alpha_2\beta|ac - a|\alpha_2\gamma|ac - a|\alpha\beta_2|bc - 1|\alpha_2\gamma|bac \\ &\quad - 1|\alpha\beta_2|abc, \\ h_2^\rho(1|\alpha\gamma|1) &= -abc|\alpha_3|1 - 2aba|\alpha_2\beta|1 - 3ab|\alpha_3|b - ab|\beta_3|c - 2bc|\beta_3|c \\ &\quad - 2ba|\alpha_2\gamma|c - ba|\alpha\beta_2|a + 2a|\alpha_3|ba - c|\alpha_3|bc - b|\beta_3|ac - b|\gamma_3|ab \\ &\quad - a|\alpha_2\beta|ab - b|\alpha_2\gamma|bc + c|\alpha_2\gamma|ac - c|\alpha\beta_2|ab - 1|\alpha_2\beta|bac \\ &\quad - 2|\alpha_2\gamma|abc. \end{aligned}$$

Using the previous results together with Theorem 2.7 we obtain the Gerstenhaber bracket between  $X_i$  for  $i \in \llbracket 1, 14 \rrbracket$  and  $X_1, X_2, X_3$ .

PROPOSITION 4.4. *The Gerstenhaber bracket on  $\mathrm{HH}^\bullet(A)$  of  $X_i$  for  $i \in \llbracket 1, 14 \rrbracket$  with an element  $X_j$  for  $j \in \llbracket 1, 3 \rrbracket$  is given by*

$$[X_i, X_1] = \begin{cases} -2X_1, & \text{if } i = 8, \\ -4X_1(X_9 + X_{10}), & \text{if } i = 13, \\ 0, & \text{if } i \in \llbracket 1, 14 \rrbracket \setminus \{8, 13\}, \end{cases}$$

$$[X_i, X_2] = \begin{cases} -2X_2, & \text{if } i = 8, \\ -4X_1X_{10}, & \text{if } i = 13, \\ 0, & \text{if } i \in \llbracket 1, 14 \rrbracket \setminus \{8, 13\}, \end{cases}$$

and

$$[X_i, X_3] = \begin{cases} 0, & \text{if } i \in \llbracket 1, 7 \rrbracket, \\ -4X_3, & \text{if } i = 8, \\ -2X_{i-5}, & \text{if } i = 9, 10, \\ 2X_6, & \text{if } i = 11, \\ 2X_7 - X_1X_8 + X_2X_8, & \text{if } i = 12, \\ -4X_3(X_9 + X_{10} + X_{11}), & \text{if } i = 13, \\ X_{13} - (2/3)X_8(X_9 + X_{10} + X_{11}), & \text{if } i = 14. \end{cases}$$

*Proof.* Note that  $\ell_{ab+ba}i_0 = Y_1$ ,  $\ell_{ab+bc-ac}i_0 = Y_2$  and  $\ell_{abac}i_0 = Y_3$ . Applying Theorem 2.7 together with Facts 4.1, 4.2 and 4.3, we get the brackets

$$[X_i, X_1] = \begin{cases} -2X_1, & \text{if } i = 8, \\ -(\alpha_2 + \beta_2 + \gamma_2)|(ab + ba) - \alpha\beta|ba - \alpha\gamma|ab, & \text{if } i = 13, \\ 0, & \text{if } i \in \llbracket 1, 12 \rrbracket \setminus \{8\}, \end{cases}$$

$$[X_i, X_2] = \begin{cases} -2X_2, & \text{if } i = 8, \\ (\alpha_2 + \beta_2 + \gamma_2)|(ac - ab - bc) + \alpha\beta|ac - \alpha\gamma|(ab + bc), & \text{if } i = 13, \\ 0, & \text{if } i \in \llbracket 1, 12 \rrbracket \setminus \{8\}, \end{cases}$$

and

$$[X_i, X_3] = \begin{cases} 0, & \text{if } i \in \llbracket 1, 7 \rrbracket, \\ -4X_3, & \text{if } i = 8, \\ -2X_{i-5}, & \text{if } i = 9, 10, \\ 2X_6, & \text{if } i = 11, \\ \alpha|(aba - abc) - \beta|bac - \gamma|bac, & \text{if } i = 12, \\ -4(\alpha_2 + \beta_2 + \gamma_2)|abac, & \text{if } i = 13. \end{cases}$$

Indeed, this was simply done by computing  $[Y_i, Y_1] = Y_i h_{\mathfrak{h}(Y_i)-1}^{ab+ba}$ ,  $[Y_i, Y_2] = Y_i h_{\mathfrak{h}(Y_i)-1}^{ab+bc-ac}$ , and  $[Y_i, Y_3] = Y_i h_{\mathfrak{h}(Y_i)-1}^{abac}$ , where  $\mathfrak{h}(Y_i)$  denotes the cohomological



degree of  $Y_i$  for  $i \in \llbracket 1, 13 \rrbracket$ , and by transport of structures. Note that the vanishing of  $[X_i, X_3]$  for  $i \in \llbracket 4, 7 \rrbracket$  also follows from a simple degree argument using Corollary 2.5 together with [3], Cor. 5.9. The latter two results also tell us that  $[X_{14}, X_j] = 0$  (or  $[Y_{14}, Y_j] = 0$ ) for  $j = 1, 2$ , by degree reasons. This result also follows from noting that  $h_3^{ab+ba}$  is of internal degree 2, so  $h_3^{ab+ba}(1|u|1)$  is of internal degree 5 for any  $u \in \mathfrak{B}_3^{!*}$ , which implies that  $Y_{14}(h_3^{ab+ba}(1|u|1))$  vanishes, since  $Y_{14}$  vanishes on any homogeneous element of internal degree strictly less than 6. Hence,  $Y_{14}h_3^{ab+ba} = 0$ . We get  $Y_{14}h_3^{ab+bc-ac} = 0$  for the same reason.

Next, we compute  $\varphi = [Y_{14}, Y_3] = Y_{14}h_3^{abc}$ . By (2.3), the map  $h_3^{abc} : P_3^b \rightarrow P_4^b$  satisfies  $\delta_4^b h_3^{abc} = \eta_3 - h_2^{abc} \delta_3^b$ . It is easy to check that

$$\begin{aligned} (\eta_3 - h_2^{abc} \delta_3^b)(1|\alpha_3|1) &= -bac|\alpha_3|a + abc|\beta_3|a - aba|\beta_3|c + aba|\alpha_2\gamma|a + v_{\alpha_3}, \\ (\eta_3 - h_2^{abc} \delta_3^b)(1|\beta_3|1) &= -2aba|\alpha_3|c + bac|\alpha_3|b + aba|\beta_3|a - abc|\beta_3|b \\ &\quad + aba|\alpha_2\gamma|b + v_{\beta_3}, \\ (\eta_3 - h_2^{abc} \delta_3^b)(1|\gamma_3|1) &= abc|\beta_3|c + 3aba|\gamma_3|c - bac|\alpha\beta_2|c + v_{\gamma_3}, \\ (\eta_3 - h_2^{abc} \delta_3^b)(1|\alpha_2\beta|1) &= abc|\alpha_3|a - 4bac|\alpha_3|b - 2aba|\beta_3|a - abc|\beta_3|b \\ &\quad + bac|\beta_3|c + 3aba|\gamma_3|b + 2aba|\alpha_2\beta|a - aba|\alpha_2\gamma|b \\ &\quad - 2abc|\alpha_2\gamma|c - abc|\alpha\beta_2|a + aba|\alpha\beta_2|c + v_{\alpha_2\beta}, \quad (4.1) \\ (\eta_3 - h_2^{abc} \delta_3^b)(1|\alpha_2\gamma|1) &= -4bac|\alpha_3|c + bac|\beta_3|a - 4abc|\beta_3|c + 2aba|\alpha_2\beta|b \\ &\quad + bac|\alpha_2\gamma|b - 3aba|\alpha_2\gamma|c + aba|\alpha\beta_2|a + v_{\alpha_2\gamma}, \\ (\eta_3 - h_2^{abc} \delta_3^b)(1|\alpha\beta_2|1) &= -3aba|\alpha_3|b + 2abc|\alpha_3|c - 4abc|\beta_3|a + bac|\beta_3|b \\ &\quad + 3aba|\gamma_3|a + 2aba|\alpha_2\beta|c - aba|\alpha_2\gamma|a - abc|\alpha_2\gamma|b \\ &\quad - bac|\alpha_2\gamma|c + 2aba|\alpha\beta_2|b + v_{\alpha\beta_2}, \end{aligned}$$

where  $v_u \in \bigoplus_{j \in \llbracket 0, 4 \rrbracket \setminus \{3\}} (A_j \otimes (A_{-3}^1)^* \otimes A_{4-j})$  for  $u \in \mathfrak{B}_3^{!*}$ . By degree reasons, the element  $h_3^{abc}(1|u|1)$  for  $u \in \mathfrak{B}_3^{!*}$  is of the form

$$\begin{aligned} h_3^{abc}(1|u|1) &= B_u + \omega_1(\lambda_1^u a|\epsilon^1|1 + \lambda_2^u 1|\epsilon^1|a + \lambda_3^u b|\epsilon^1|1 + \lambda_4^u 1|\epsilon^1|b + \lambda_5^u c|\epsilon^1|1 \\ &\quad + \lambda_6^u 1|\epsilon^1|c), \end{aligned}$$

where  $B_u \in K_4^b$  has internal degree 7 and  $\lambda_i^u \in \mathbb{k}$  for  $i \in \llbracket 1, 6 \rrbracket$ . Therefore,

$$\begin{aligned} (\eta_3 - h_2^{abc} \delta_3^b)(1|u|1) &= \delta_4^b h_3^{abc}(1|u|1) \\ &= d_4^b(B_u) + \lambda_1^u f_0^b(a|\epsilon^1|1) + \lambda_2^u f_0^b(1|\epsilon^1|a) + \lambda_3^u f_0^b(b|\epsilon^1|1) \\ &\quad + \lambda_4^u f_0^b(1|\epsilon^1|b) + \lambda_5^u f_0^b(c|\epsilon^1|1) + \lambda_6^u f_0^b(1|\epsilon^1|c). \end{aligned} \quad (4.2)$$

Using the explicit expression of the differential  $d_4^b$  given in [3], Fact 3.1, together with an elementary computation we see that, given any homogeneous element  $B \in K_4^b$  of internal degree 7, the coefficients of  $aba|\gamma_3|a$  and  $aba|\alpha_2\gamma|a$  in  $d_4^b(B)$

are equal, the coefficients of  $aba|\gamma_3|b$  and  $aba|\alpha_2\gamma|b$  in  $d_4^b(B)$  coincide, and the coefficients of  $abc|\beta_3|c$  and  $abc|\alpha_2\beta|c$  in  $d_4^b(B)$  are also the same. Comparing the coefficients of  $aba|\gamma_3|a$  and  $aba|\alpha_2\gamma|a$  in both sides of the equation (4.2), where the left member is explicitly given by (4.1) and the right member is computed using Fact 3.2, we obtain

$$\lambda_1^{\alpha_3} + \lambda_2^{\alpha_3} = 1/3, \quad \lambda_1^{\alpha\beta_2} + \lambda_2^{\alpha\beta_2} = -4/3, \quad \lambda_1^u + \lambda_2^u = 0$$

for  $u \in \mathfrak{B}_3^{!*} \setminus \{\alpha_3, \alpha\beta_2\}$ . Similarly, comparing the coefficients of  $aba|\gamma_3|b$  and  $aba|\alpha_2\gamma|b$  in both sides of the equation (4.2), where the left member is explicitly given by (4.1) and the right member is computed using Fact 3.2, we obtain

$$\lambda_3^{\beta_3} + \lambda_4^{\beta_3} = 1/3, \quad \lambda_3^{\alpha_2\beta} + \lambda_4^{\alpha_2\beta} = -4/3, \quad \lambda_3^u + \lambda_4^u = 0$$

for  $u \in \mathfrak{B}_3^{!*} \setminus \{\beta_3, \alpha_2\beta\}$ . Comparing the coefficients of  $abc|\beta_3|c$  and  $abc|\alpha_2\beta|c$  in both sides of the equation (4.2), where the left member is explicitly given by (4.1) and the right member is computed using Fact 3.2, we obtain

$$\lambda_5^{\gamma_3} + \lambda_6^{\gamma_3} = 1/3, \quad \lambda_5^{\alpha_2\gamma} + \lambda_6^{\alpha_2\gamma} = -4/3, \quad \lambda_5^u + \lambda_6^u = 0$$

for  $u \in \mathfrak{B}_3^{!*} \setminus \{\gamma_3, \alpha_2\gamma\}$ . Then  $\varphi(1|u|1) = Y_{14}h_3^{abc}(1|u|1)$  for  $u \in \mathfrak{B}_3^{!*}$  is given by

$$\begin{aligned} \varphi(1|\alpha_3|1) &= (1/3)a, & \varphi(1|\beta_3|1) &= (1/3)b, & \varphi(1|\gamma_3|1) &= (1/3)c, \\ \varphi(1|\alpha_2\beta|1) &= -(4/3)b, & \varphi(1|\alpha_2\gamma|1) &= -(4/3)c, & \varphi(1|\alpha\beta_2|1) &= -(4/3)a. \end{aligned}$$

Hence,  $[X_{14}, X_3] = (1/3)(\alpha_3|a + \beta_3|b + \gamma_3|c) - (4/3)(\alpha_2\beta|b + \alpha_2\gamma|c + \alpha\beta_2|a)$ . We now note the following identities,

$$\begin{aligned} \alpha_2|(ab + ba) &= X_1X_9, & \beta_2|(ab + ba) &= X_1X_{10}, \\ \alpha|aba + \beta|bac &= (1/2)(X_1X_8 - X_2X_8), & \alpha_2|abac &= X_3X_9, \\ \beta_2|abac &= X_3X_{10}, & \gamma_2|abac &= X_3X_{11} \end{aligned} \tag{4.3}$$

and

$$\begin{aligned} (\alpha_3 - \alpha\beta_2)|a &= (1/2)\{X_{13} + X_8(X_9 - X_{10} - X_{11})\}, \\ (\beta_3 - \alpha_2\beta)|b &= (1/2)\{X_{13} + X_8(X_{10} - X_9 - X_{11})\}, \\ (\gamma_3 - \alpha_2\gamma)|c &= (1/2)\{X_{13} + X_8(X_{11} - X_9 - X_{10})\}, \end{aligned}$$

given in [3], (6.2). Using the previous equalities as well as the coboundaries  $g_{j,2}^2 \in \tilde{\mathfrak{B}}_2^2$  for  $j \in \llbracket 1, 8 \rrbracket \setminus \{4, 5\}$  and  $e_{1,3}^1 = \alpha|(aba + abc) + (\beta - \gamma)|bac \in \tilde{\mathfrak{B}}_3^1$  of the sets  $\tilde{\mathfrak{B}}_2^2$  and  $\tilde{\mathfrak{B}}_3^1$  given in [3], Subsubsection 5.3.1, we get

$$\begin{aligned} [X_{13}, X_1] &= -(\alpha_2 + \beta_2 + \gamma_2)|(ab + ba) - \alpha\beta|ba - \alpha\gamma|ab - 3g_{1,2}^2 - 3g_{2,2}^2 - 2g_{3,2}^2 \\ &\quad + g_{8,2}^2 \\ &= -4(\alpha_2 + \beta_2)|(ab + ba) = -4X_1(X_9 + X_{10}), \end{aligned}$$

$$\begin{aligned}
 [X_{13}, X_2] &= (\alpha_2 + \beta_2 + \gamma_2)(ac - ab - bc) + \alpha\beta|ac - \alpha\gamma|(ab + bc) - g_{1,2}^2 \\
 &\quad - 2g_{2,2}^2 - g_{3,2}^2 + g_{6,2}^2 - g_{7,2}^2 + g_{8,2}^2 \\
 &= -4\beta_2(ab + ba) = -4X_1X_{10}, \\
 [X_{12}, X_3] &= \alpha|(aba - abc) - \beta|bac - \gamma|bac - e_{1,3}^1 \\
 &= 2\alpha|(aba - abc) - 2(\alpha|aba + \beta|bac) = 2X_7 - X_1X_8 + X_2X_8, \\
 [X_{13}, X_3] &= -4(\alpha_2 + \beta_2 + \gamma_2)|abac = -4X_3(X_9 + X_{10} + X_{11}), \\
 [X_{14}, X_3] &= (1/3)(\alpha_3|a + \beta_3|b + \gamma_3|c) - (4/3)(\alpha_2\beta|b + \alpha_2\gamma|c + \alpha\beta_2|a) \\
 &= X_{13} - (2/3)X_8(X_9 + X_{10} + X_{11}).
 \end{aligned}$$

The proposition is thus proved. □

#### 4.2 GERSTENHABER BRACKETS OF $\mathrm{HH}^1(A)$ WITH $\mathrm{HH}^n(A)$

In this subsection we are going to utilize the method recalled in Subsection 2.3 to compute the Gerstenhaber bracket of  $X_i$  for  $i \in \llbracket 4, 8 \rrbracket$  with the elements  $X_j$  for  $j \in \llbracket 1, 14 \rrbracket$ .

Let  $\rho : A \rightarrow A$  be a derivation of  $A$ . By [8], Lemma 1.3, the  $\rho^\epsilon$ -lifting  $\rho_\bullet = \{\rho_n : P_n^b \rightarrow P_n^b\}_{n \in \mathbb{N}_0}$  of  $\rho$  to  $(P_\bullet^b, \delta_\bullet^b)$  exists, and it can be chosen in such a way that

$$\begin{aligned}
 \rho_0(x|\epsilon^1|y) &= \rho(x)|\epsilon^1|y + x|\epsilon^1|\rho(y), \\
 \rho_n(\omega_i x|u|y) &= xq_{\omega_i u}y + \omega_i \rho(x)|u|y + \omega_i x|u|\rho(y),
 \end{aligned} \tag{4.4}$$

for all  $x, y \in A$ ,  $n \in \mathbb{N}$ ,  $i \in \llbracket 0, \lfloor n/4 \rfloor \rrbracket$  and  $u \in \mathcal{B}_{n-4i}^{1*}$ , where  $q_{\omega_i u} \in P_n^b$  satisfies that  $\delta_n^b(q_{\omega_i u}) = \rho_{n-1} \delta_n^b(\omega_i 1|u|1)$ . To reduce space, we will usually write  $q_u$  instead of  $q_{\omega_0 u}$ . As recalled in Subsection 2.3, given  $\phi \in \mathrm{HH}^n(A)$ , the Gerstenhaber bracket  $[G(\rho)i_1, \phi] \in \mathrm{HH}^n(A)$  is given by the cohomology class of  $\rho\phi - \phi\rho_n$ .

In what follows, we consider a set of derivations of  $A$  whose classes give a basis of  $\mathrm{HH}^1(A)$  and for each of them we will provide some of the corresponding elements  $q_{\omega_i u}$  satisfying (4.4). Then, we shall compute the respective Gerstenhaber brackets by means of Theorem 2.13.

The proof of the following result follows immediately from the statement.

**PROPOSITION 4.5.** *Let  $\rho : A \rightarrow A$  be the derivation of  $A$  defined by  $\rho(x) = \deg(x)x$  for  $x \in \mathcal{B}$ . Then  $\rho_\bullet$  defined by  $\rho_n(\omega_i x|u|y) = (\deg(x) + \deg(y) + n + 2i)\omega_i x|u|y$  for  $x, y \in \mathcal{B}$ ,  $i \in \llbracket 0, \lfloor n/4 \rfloor \rrbracket$ ,  $u \in \mathcal{B}_{n-4i}^{1*}$  and  $n \in \mathbb{N}_0$  is a  $\rho^\epsilon$ -lifting of  $\rho$ . Note that  $\deg(x) + \deg(y) + n + 2i$  is the internal degree of  $\omega_i x|u|y$ . Since  $G(\rho)i_1 = -X_8$ , the Gerstenhaber bracket  $[X_8, \phi] \in \mathrm{HH}^n(A)$  for  $\phi \in \mathrm{HH}^n(A)$  is given by the cohomology class  $-\mathfrak{a}(\phi)\phi$ , where  $\mathfrak{a}(\phi)$  is the internal degree of  $\phi$ .*

Hence,

$$[X_8, X_j] = \begin{cases} -2X_j, & \text{if } j \in [1, 7] \setminus \{3\}, \\ -4X_3, & \text{if } j = 3, \\ 0, & \text{if } j = 8, \\ 2X_j, & \text{if } j \in [9, 13], \\ 6X_{14}, & \text{if } j = 14. \end{cases}$$

The proof of Facts 4.6, 4.7, 4.8 and 4.9 below is a lengthy but straightforward computation.

FACT 4.6. Let  $\rho = \rho^4 : A \rightarrow A$  be the derivation of  $A$  defined by  $\rho^4(a) = bac$  and  $\rho^4(x) = 0$  for  $x \in \mathfrak{B} \setminus \{a\}$ . Then the elements  $q_{\omega_i u} = q_{\omega_i u}^4 \in P_n^b$  in (4.4) can be chosen as follows. First,  $q_{\beta_n}^4 = q_{\gamma_n}^4 = 0$  for  $n \in \mathbb{N}$ . Moreover,

$$\begin{aligned} q_\alpha^4 &= ba|\gamma|1 + b|\alpha|c + 1|\beta|ac, \\ q_{\alpha_2}^4 &= ba|\alpha\beta|1 - b|\alpha_2|b - c|\alpha_2|c - b|\alpha\beta|c + 1|\alpha\beta|ac, \\ q_{\alpha\beta}^4 &= ab|\gamma_2|1 - ab|\alpha_2|1 + ba|\alpha\gamma|1 - 2b|\alpha_2|c - c|\alpha_2|a - b|\beta_2|c - a|\alpha\beta|c \\ &\quad - b|\alpha\beta|a + 1|\alpha_2|bc - 1|\beta_2|ab + 1|\alpha\gamma|ac, \\ q_{\alpha\gamma}^4 &= ba|\gamma_2|1 + 1|\beta_2|ac, \\ q_{\alpha_3}^4 &= bc|\beta_3|1 + ba|\alpha_2\gamma|1 + b|\alpha_3|c + c|\alpha_3|b + b|\beta_3|a - c|\beta_3|c - a|\alpha_2\gamma|c \\ &\quad + b|\alpha_2\gamma|b - 1|\beta_3|ac - 1|\alpha_2\gamma|bc, \\ q_{\alpha_2\beta}^4 &= ab|\gamma_3|1 + ba|\alpha\beta_2|1 - a|\alpha_3|b - 2a|\beta_3|c - c|\beta_3|a - b|\gamma_3|c - c|\gamma_3|b \\ &\quad - a|\alpha_2\beta|c - c|\alpha\beta_2|c - b|\alpha_2\gamma|c - a|\alpha_2\gamma|a + 1|\alpha_3|bc + 1|\gamma_3|ba - 1|\alpha_2\gamma|ba \\ &\quad - 1|\alpha\beta_2|bc, \\ q_{\alpha_2\gamma}^4 &= ba|\alpha_2\beta|1 + ab|\alpha\beta_2|1 - 2ab|\alpha_3|1 - ba|\beta_3|1 + a|\alpha_3|c + 2b|\alpha_3|b + c|\alpha_3|a \\ &\quad + a|\beta_3|a + b|\beta_3|c + c|\beta_3|b + a|\alpha_2\gamma|b + a|\alpha\beta_2|c - 1|\alpha_3|ba - 2|\beta_3|ab \\ &\quad + 1|\alpha\beta_2|ac, \\ q_{\alpha\beta_2}^4 &= 3ba|\gamma_3|1 - (ab + bc)|\beta_3|1 + (ab + bc)|\alpha_2\beta|1 - 3c|\alpha_3|b - 2a|\beta_3|b - b|\beta_3|a \\ &\quad - c|\beta_3|c - 2a|\gamma_3|c - 2c|\gamma_3|a - a|\alpha_2\beta|b - 2c|\alpha_2\beta|c - 2b|\alpha_2\gamma|b - c|\alpha_2\gamma|a \\ &\quad - c|\alpha\beta_2|b + 2|\beta_3|ac + 1|\gamma_3|ab - 1|\alpha_2\gamma|ab, \\ q_{\omega_1 \epsilon^1}^4 &= bac|\alpha_4|a + 4abc|\beta_4|b - aba|\gamma_4|c + 4ab|\alpha_3\beta|bc - 4bc|\alpha_3\beta|ab \\ &\quad + 2(ba + ac)|\alpha_3\gamma|ac + 2ba|\alpha_3\gamma|ba + ab|\alpha_2\beta_2|ac + bc|\alpha_2\beta_2|ba \\ &\quad - 2ba|\alpha_2\beta_2|(ab + bc) - 4ab|\alpha_4|ba - 2bc|\alpha_4|ba + (ba + ac)|\alpha_4|bc \\ &\quad + 4ab|\alpha_4|ac - 2(ba + ac)|\alpha_4|ab + 8bc|\beta_4|ac - 10ba|\beta_4|ab - 2ac|\beta_4|ab \\ &\quad + 6ab|\beta_4|ac + 4(ab + bc)|\beta_4|ba - 2ab|\gamma_4|ba - 4bc|\gamma_4|ba + 6ab|\gamma_4|ac \\ &\quad - 5ba|\gamma_4|ab + 4ba|\gamma_4|bc - 6ac|\gamma_4|ab + a|\alpha_4|bac + 4b|\beta_4|abc - c|\gamma_4|aba \\ &\quad - \omega_1 c|\epsilon^1|c. \end{aligned}$$

FACT 4.7. Let  $\rho = \rho^5 : A \rightarrow A$  be the derivation of  $A$  defined by  $\rho^5(b) = abc$  and  $\rho^5(x) = 0$  for  $x \in \mathfrak{B} \setminus \{b\}$ . Then the elements  $q_{\omega_i u} = q_{\omega_i u}^5 \in P_n^b$  in (4.4) can be chosen as follows. First,  $q_{\alpha_n}^5 = q_{\gamma_n}^5 = 0$  for  $n \in \mathbb{N}$ . Moreover,

$$\begin{aligned} q_\beta^5 &= ab|\gamma|1 + a|\beta|c + 1|\alpha|bc, \\ q_{\beta_2}^5 &= ab|\alpha\gamma|1 - a|\beta_2|a - c|\beta_2|c - a|\alpha\gamma|c + 1|\alpha\gamma|bc, \\ q_{\alpha\beta}^5 &= ab|\gamma_2|1 + 1|\alpha_2|bc, \\ q_{\alpha\gamma}^5 &= ab|\alpha\beta|1 - bc|\alpha\beta|1 - 2ba|\beta_2|1 - a|\alpha_2|c - 2a|\beta_2|c + b|\beta_2|a - c|\beta_2|b \\ &\quad + b|\gamma_2|a + b|\alpha\beta|b - a|\alpha\gamma|b + 1|\beta_2|ac - 1|\alpha_2|ba + 1|\alpha\beta|bc, \\ q_{\beta_3}^5 &= (ba + ac)|\alpha_3|1 + ab|\alpha_2\gamma|1 + a|\alpha_3|b + b|\alpha_3|a + 2a|\beta_3|c + c|\beta_3|a + b|\gamma_3|c \\ &\quad + c|\gamma_3|b + a|\alpha_2\beta|c + a|\alpha_2\gamma|a + c|\alpha\beta_2|c - 1|\gamma_3|ba + 1|\alpha\beta_2|bc, \\ q_{\alpha_2\beta}^5 &= 2ab|\gamma_3|1 - 2(ba + ac)|\alpha_3|1 - (ab + bc)|\alpha_2\gamma|1 + (ba + ac)|\alpha\beta_2|1 - a|\alpha_3|b \\ &\quad - 2b|\alpha_3|a - 2c|\beta_3|a - b|\gamma_3|c - c|\gamma_3|b - 2a|\alpha_2\gamma|a - c|\alpha\beta_2|c + 2|\alpha_3|bc \\ &\quad + 1|\gamma_3|ba - 1|\alpha_2\gamma|ba, \\ q_{\alpha_2\gamma}^5 &= ba|\alpha_2\beta|1 - 2ba|\beta_3|1 - ab|\alpha_3|1 + a|\alpha_3|c + b|\alpha_3|b + c|\alpha_3|a + 2a|\beta_3|a \\ &\quad + b|\beta_3|c + c|\beta_3|b + b|\alpha_2\beta|c + b|\alpha_2\gamma|a + b|\alpha\beta_2|b - 2|\alpha_3|ba - 1|\beta_3|ab \\ &\quad + 1|\alpha_2\beta|bc, \\ q_{\alpha\beta_2}^5 &= ba|\gamma_3|1 + ab|\alpha_2\beta|1 - b|\alpha_3|c - c|\alpha_3|b - b|\beta_3|a + c|\beta_3|c - b|\alpha_2\gamma|b \\ &\quad + b|\alpha\beta_2|c + 2|\beta_3|ac - 1|\alpha_2\gamma|ab + 1|\alpha_2\gamma|bc, \\ q_{\omega_1\epsilon^!}^5 &= 4bac|\alpha_4|a + abc|\beta_4|b - aba|\gamma_4|c + ab|\alpha_3\beta|ab + 5ab|\alpha_3\beta|bc \\ &\quad - 4bc|\alpha_3\beta|ab + 2(ba + ac)|\alpha_3\gamma|ac + 2ba|\alpha_3\gamma|ba - ac|\alpha_3\gamma|ba \\ &\quad - ab|\alpha_2\beta_2|ba + 2bc|\alpha_2\beta_2|ba - 2ba|\alpha_2\beta_2|ab - ba|\alpha_2\beta_2|bc + ac|\alpha_2\beta_2|bc \\ &\quad - 11ab|\alpha_4|ba + 2ab|\alpha_4|ac + 6ba|\alpha_4|bc + ac|\alpha_4|ab + 9ac|\alpha_4|bc \\ &\quad + 3ab|\beta_4|ac + bc|\beta_4|ac - 6ba|\beta_4|ab + ba|\beta_4|bc - 3ac|\beta_4|ab - 5ab|\gamma_4|ba \\ &\quad + 4ab|\gamma_4|ac - 4bc|\gamma_4|ba - 2bc|\gamma_4|ac - 3ba|\gamma_4|ab + 6ba|\gamma_4|bc - 8ac|\gamma_4|ab \\ &\quad + 4a|\alpha_4|bac + b|\beta_4|abc - c|\gamma_4|aba - \omega_1c|\epsilon^!|c. \end{aligned}$$

FACT 4.8. Let  $\rho = \rho^6 : A \rightarrow A$  be the derivation of  $A$  defined by  $\rho^6(c) = aba$  and  $\rho^6(x) = 0$  for  $x \in \mathfrak{B} \setminus \{c\}$ . Then the elements  $q_{\omega_i u} = q_{\omega_i u}^6 \in P_n^b$  in (4.4) can be chosen as follows. First,  $q_{\alpha_n}^6 = q_{\beta_n}^6 = 0$  for  $n \in \mathbb{N}$ . Moreover,

$$\begin{aligned} q_\gamma^6 &= ab|\alpha|1 + a|\beta|a + 1|\alpha|ba, \\ q_{\gamma_2}^6 &= ba|\alpha\beta|1 - b|\alpha_2|b + c|\beta_2|c + a|\gamma_2|a + c|\alpha\beta|a + a|\alpha\gamma|c + 1|\alpha\gamma|ab, \\ q_{\alpha\beta}^6 &= 2ab|\alpha_2|1 + c|\alpha_2|a + b|\beta_2|c + b|\alpha\beta|a + 1|\beta_2|ab, \\ q_{\alpha\gamma}^6 &= ba|\beta_2|1 + a|\alpha_2|c + c|\beta_2|b + a|\alpha\gamma|b + 2|\alpha_2|ba, \\ q_{\gamma_3}^6 &= ab|\alpha\beta_2|1 - ba|\beta_3|1 + a|\beta_3|a - a|\gamma_3|b - b|\gamma_3|a - b|\alpha_2\beta|c - c|\alpha_2\beta|b \\ &\quad - b|\alpha\beta_2|b - 1|\beta_3|ab + 1|\alpha\beta_2|ba, \end{aligned}$$

$$\begin{aligned}
q_{\alpha_2\beta}^6 &= ac|\alpha_3|1 + ab|\alpha_2\gamma|1 + 2c|\alpha_3|c + 2a|\beta_3|c + 2c|\beta_3|a + 2a|\alpha_2\gamma|a + b|\alpha_2\gamma|c \\
&\quad + c|\alpha_2\gamma|b + a|\alpha\beta_2|b + b|\alpha\beta_2|a - 1|\alpha_3|(ab + bc) + 1|\alpha_2\gamma|ba, \\
q_{\alpha_2\gamma}^6 &= 3ab|\alpha_3|1 + 2ba|\beta_3|1 - a|\alpha_3|c - c|\alpha_3|a - b|\beta_3|c - c|\beta_3|b + 3|\alpha_3|ba \\
&\quad + 2|\beta_3|ab, \\
q_{\alpha\beta_2}^6 &= 2bc|\beta_3|1 + 2ba|\alpha_2\gamma|1 + 3b|\alpha_3|c + 3c|\alpha_3|b + 3b|\alpha_2\gamma|b - 2|\beta_3|(ba + ac) \\
&\quad + 2|\alpha_2\gamma|ab, \\
q_{\omega_1\epsilon^1}^6 &= 2bac|\alpha_2\beta_2|a + aba|\alpha_2\beta_2|c - 5bac|\alpha_4|a - 3abc|\beta_4|b - 2bac|\alpha_3\beta|b \\
&\quad + 8ab|\alpha_4|ba - 6ab|\alpha_4|ac + 6bc|\alpha_4|ba + 3ba|\alpha_4|bc + 3ac|\alpha_4|bc \\
&\quad + 3ab|\beta_4|ac + 3bc|\beta_4|ac + 10ba|\beta_4|ab - 8ba|\beta_4|bc + 8ac|\beta_4|ab \\
&\quad - 4ab|\gamma_4|ba - 2ab|\alpha_2\beta_2|ba - 5a|\alpha_4|bac - 3b|\beta_4|abc - 2b|\alpha_3\gamma|bac \\
&\quad + 2a|\alpha_2\beta_2|bac + c|\alpha_2\beta_2|aba - \omega_1a|\epsilon^1|a.
\end{aligned}$$

FACT 4.9. Let  $\rho = \rho^7 : A \rightarrow A$  be the derivation of  $A$  defined by  $\rho^7(a) = aba - abc$ ,  $\rho^7(ab) = \rho^7(ac) = abac$ ,  $\rho^7(ba) = -abac$  and  $\rho^7(x) = 0$  for  $x \in \mathfrak{B} \setminus \{a, ab, ba, ac\}$ . Then the elements  $q_{\omega_i u} = q_{\omega_i u}^7 \in P_n^b$  in (4.4) can be chosen as follows. First,  $q_{\beta_n}^7 = q_{\gamma_n}^7 = 0$  for  $n \in \mathbb{N}$ . Moreover,

$$\begin{aligned}
q_{\alpha}^7 &= ab|\alpha|1 - ab|\gamma|1 + a|\beta|a - a|\beta|c + 1|\alpha|ba - 1|\alpha|bc, \\
q_{\alpha_2}^7 &= ab|\alpha_2|1 + ac|\alpha_2|1 - a|\alpha\beta|a - 1|\alpha_2|bc + 2|\alpha_2|ba, \\
q_{\alpha\beta}^7 &= ba|\beta_2|1 - ba|\gamma_2|1 + (ba + ac)|\alpha\beta|1 + a|\alpha_2|c + c|\beta_2|b + c|\beta_2|c - a|\gamma_2|c \\
&\quad + b|\gamma_2|b - c|\gamma_2|b + c|\alpha\beta|a - c|\alpha\beta|c + a|\alpha\gamma|b + c|\alpha\gamma|b + 1|\alpha_2|ba \\
&\quad - 1|\gamma_2|ba + 1|\alpha\beta|(ba + ac), \\
q_{\alpha\gamma}^7 &= ab|\alpha_2|1 - ab|\gamma_2|1 + c|\alpha_2|a - c|\alpha_2|c + b|\beta_2|c + b|\alpha\beta|a - b|\alpha\beta|c + 1|\beta_2|ab \\
&\quad + 1|\alpha\beta|ac, \\
q_{\alpha_3}^7 &= ab|\alpha_3|1 + ac|\alpha_3|1 - 1|\alpha_3|bc + 2|\alpha_3|ba, \\
q_{\alpha_2\beta}^7 &= ab|\alpha_3|1 + 2ba|\beta_3|1 + 2bc|\beta_3|1 - ba|\gamma_3|1 + 2ba|\alpha_2\gamma|1 - ab|\alpha\beta_2|1 - a|\alpha_3|c \\
&\quad + 3b|\alpha_3|c + c|\alpha_3|b - a|\beta_3|a - a|\beta_3|b + b|\beta_3|a - b|\beta_3|c - 2c|\beta_3|b + c|\beta_3|c \\
&\quad - a|\gamma_3|b + a|\gamma_3|c - b|\gamma_3|b + 2c|\gamma_3|a + c|\alpha_2\beta|c - a|\alpha_2\gamma|b + a|\alpha_2\gamma|c \\
&\quad + b|\alpha_2\gamma|b - a|\alpha\beta_2|c + b|\alpha\beta_2|c - c|\alpha\beta_2|b + 2|\alpha_3|ba + 1|\beta_3|ab - 3|\gamma_3|ab \\
&\quad - 1|\alpha_2\beta|bc + 1|\alpha_2\beta|ba + 2|\alpha_2\beta|ac, \\
q_{\alpha_2\gamma}^7 &= ac|\alpha_3|1 - ba|\gamma_3|1 - (ab + bc)|\alpha_2\beta|1 + ba|\alpha\beta_2|1 - b|\alpha_3|a + 3b|\alpha_3|c \\
&\quad + 2c|\alpha_3|b + c|\alpha_3|c + a|\beta_3|c + c|\beta_3|a - a|\gamma_3|a + 2a|\gamma_3|c - 2b|\gamma_3|b \\
&\quad + 2c|\gamma_3|a + b|\alpha_2\beta|b - c|\alpha_2\beta|a + 2c|\alpha_2\beta|c + a|\alpha_2\gamma|a + b|\alpha_2\gamma|b + b|\alpha_2\gamma|c \\
&\quad + c|\alpha_2\gamma|a + b|\alpha\beta_2|a - 2|\gamma_3|ab + 2|\alpha_2\beta|ac, \\
q_{\alpha\beta_2}^7 &= ab|\alpha_3|1 + ba|\beta_3|1 - 2ab|\gamma_3|1 - a|\alpha_3|c - c|\alpha_3|a + 2c|\alpha_3|c - a|\gamma_3|b \\
&\quad + b|\gamma_3|c + c|\alpha_2\gamma|b + 1|\alpha_3|ba + 1|\beta_3|ab - 1|\gamma_3|ba,
\end{aligned}$$

$$\begin{aligned}
 q_{\omega_1 \epsilon^1}^7 = & 2abc|\alpha_2\beta_2|a - 3abc|\alpha_2\beta_2|c - aba|\alpha_2\beta_2|a + 2aba|\alpha_2\beta_2|b - 5bac|\alpha_4|b \\
 & - 7bac|\alpha_4|c + 5abc|\beta_4|c + 3aba|\gamma_4|a - 3aba|\gamma_4|b + 8ab|\alpha_4|ab \\
 & + 3ab|\alpha_4|bc + bc|\alpha_4|bc - 2ba|\alpha_4|ba + 4ac|\alpha_4|ba + 6ac|\alpha_4|ac + 9ab|\beta_4|ab \\
 & - 4bc|\beta_4|ab - 5bc|\beta_4|bc - 4ba|\beta_4|ba - 7ba|\beta_4|ac + 14ab|\gamma_4|ab \\
 & + 7bc|\gamma_4|ab - ba|\gamma_4|ba - ba|\gamma_4|ac - 8ac|\gamma_4|ba + 7ab|\alpha_3\beta|ac \\
 & - 3bc|\alpha_3\beta|ba + bc|\alpha_3\beta|ac + 2ba|\alpha_3\beta|ab + 3ba|\alpha_3\beta|bc + ba|\alpha_3\beta|ba \\
 & + 3ba|\alpha_3\beta|ac + 3ac|\alpha_3\beta|bc - 6ac|\alpha_3\beta|ba - 3ac|\alpha_3\beta|ac - ab|\alpha_3\gamma|ab \\
 & + 5ab|\alpha_3\gamma|bc - 9ab|\alpha_3\gamma|ba - 5ab|\alpha_3\gamma|ac + 3bc|\alpha_3\gamma|ab + 6bc|\alpha_3\gamma|bc \\
 & - 2bc|\alpha_3\gamma|ba - 2ba|\alpha_3\gamma|ab - 2ba|\alpha_3\gamma|bc + 6ac|\alpha_3\gamma|ab + 7ab|\alpha_2\beta_2|ab \\
 & + 2ab|\alpha_2\beta_2|bc - ba|\alpha_2\beta_2|ba + 2ba|\alpha_2\beta_2|ac + 4ac|\alpha_2\beta_2|ac - 2b|\alpha_4|bac \\
 & + c|\alpha_4|bac + a|\beta_4|abc - 2c|\beta_4|abc + 4a|\gamma_4|aba + 2b|\gamma_4|aba \\
 & + 2a|\alpha_2\beta_2|abc - 2c|\alpha_2\beta_2|abc - a|\alpha_2\beta_2|aba - 2b|\alpha_2\beta_2|bac - c|\alpha_2\beta_2|bac \\
 & + \omega_1 3|\epsilon^1|(ba + ac - bc).
 \end{aligned}$$

We will now apply the previous results to compute the Gerstenhaber brackets of  $X_i$  for  $i \in \llbracket 4, 7 \rrbracket$  with all the other generators of the Hochschild cohomology of  $A$ .

PROPOSITION 4.10. *The Gerstenhaber bracket  $[X_i, X_j] \in \mathrm{HH}^\bullet(A)$  for  $i \in \llbracket 4, 7 \rrbracket$  and  $j \in \llbracket 1, 14 \rrbracket$  is given by*

$$[X_i, X_j] = \begin{cases} 0, & \text{if } (i, j) \in (\llbracket 4, 7 \rrbracket \times \llbracket 1, 7 \rrbracket) \\ & \cup (\llbracket 4, 6 \rrbracket \times \llbracket 9, 11 \rrbracket), \\ 2X_i, & \text{if } i \in \llbracket 4, 7 \rrbracket \text{ and } j = 8, \\ 4X_1X_9, & \text{if } i = 7 \text{ and } j = 9, \\ X_1X_{10}, & \text{if } i = 7 \text{ and } j = 10, \\ -X_1(X_9 + X_{10}), & \text{if } i = 7 \text{ and } j = 11, \\ 2X_1X_{i+5}, & \text{if } i \in \llbracket 4, 5 \rrbracket \text{ and } j = 12, \\ 2X_1(X_9 + X_{10}), & \text{if } i = 6 \text{ and } j = 12, \\ X_1X_9, & \text{if } i = 7 \text{ and } j = 12, \\ \tau_i 8X_4X_{10}, & \text{if } i \in \llbracket 4, 6 \rrbracket \text{ and } j = 13, \\ 4X_1X_{13} - 4X_2X_{13} - 8X_4X_{12}, & \text{if } i = 7 \text{ and } j = 13, \\ \tau_i((1/3)X_{i+5}^2 - (4/3)X_9X_{10}), & \text{if } i \in \llbracket 4, 6 \rrbracket \text{ and } j = 14, \\ X_9X_{12}, & \text{if } i = 7 \text{ and } j = 14, \end{cases}$$

where  $\tau_i = 1$  if  $i \in \llbracket 4, 5 \rrbracket$  and  $\tau_6 = -1$ .

*Proof.* Given  $i \in \llbracket 4, 7 \rrbracket$ , let  $\rho^i$  be the derivation of Fact 4.6, Fact 4.7, Fact 4.8, and Fact 4.9, respectively. Note that  $G(\rho^i)i_1 = -Y_i$ . By Theorem 2.13,  $[-Y_i, Y_j]$  is precisely the cohomology class of  $\rho^i Y_j - Y_j \rho_n^i$  for  $i \in \llbracket 4, 7 \rrbracket$  and  $j \in \llbracket 1, 13 \rrbracket$ , where  $n$  is the cohomological degree of  $Y_j$  and  $\rho_n^i$  is obtained

from (4.4) together with Fact 4.6 for  $i = 4$ , Fact 4.7 for  $i = 5$ , Fact 4.8 for  $i = 6$ , and Fact 4.9 for  $i = 7$ . It is explicitly given by

$$[-X_4, X_j] = \begin{cases} 0, & \text{if } j \in \llbracket 1, 7 \rrbracket \cup \{9\}, \\ -2X_4, & \text{if } j = 8, \\ \alpha\beta|(ab + bc) - \alpha\gamma|ac, & \text{if } j = 10, \\ -\alpha\beta|ab - \alpha\gamma|ba, & \text{if } j = 11, \\ \alpha_2|(bc - ba - ac), & \text{if } j = 12, \\ 3\alpha_2\gamma|aba - 5\alpha\beta_2|bac, & \text{if } j = 13, \end{cases}$$

and

$$[-X_5, X_j] = \begin{cases} 0, & \text{if } j \in \llbracket 1, 7 \rrbracket \cup \{10\}, \\ -2X_5, & \text{if } j = 8, \\ -\alpha\beta|bc + \alpha\gamma|(ba + ac), & \text{if } j = 9, \\ -\alpha\beta|ab - \alpha\gamma|ba, & \text{if } j = 11, \\ -\beta_2|(ab + bc - ac), & \text{if } j = 12, \\ -5\alpha_2\beta|abc + 3\alpha_2\gamma|aba, & \text{if } j = 13, \end{cases}$$

as well as

$$[-X_6, X_j] = \begin{cases} 0, & \text{if } j \in \llbracket 1, 7 \rrbracket \cup \{11\}, \\ -2X_6, & \text{if } j = 8, \\ \alpha\beta|(bc - ab) - \alpha\gamma|(2ba + ac), & \text{if } j = 9, \\ -\alpha\beta|(ab + bc) + \alpha\gamma|ac, & \text{if } j = 10, \\ \gamma_2|(bc - ba - ac) - \alpha\beta|ba - \alpha\gamma|ab, & \text{if } j = 12, \\ -10\alpha_2\gamma|aba - 2\alpha\beta_2|bac, & \text{if } j = 13, \end{cases}$$

together with

$$[-X_7, X_{13}] = \alpha_3|(abc - 2aba) + \alpha_2\beta|(2bac - 6aba) - \alpha_2\gamma|(abc + 4bac) + 5\alpha\beta_2|(abc - aba),$$

and

$$[-X_7, X_j] = \begin{cases} 0, & \text{if } j \in \llbracket 1, 7 \rrbracket, \\ -2X_7, & \text{if } j = 8, \\ \alpha_2|(bc - ab - ac - 2ba) - \alpha\beta|(ba + ac) + \alpha\gamma|bc, & \text{if } j = 9, \\ \alpha\beta|ac - \alpha\gamma|(ab + bc), & \text{if } j = 10, \\ \alpha\beta|ba + \alpha\gamma|ab, & \text{if } j = 11, \\ (\alpha\beta + \alpha\gamma)|(bc - ba - ac), & \text{if } j = 12. \end{cases}$$

Next, we will compute  $\varphi^i = [-Y_i, Y_{14}] = \rho^i Y_{14} - Y_{14} \rho_4^i$  for  $i \in \llbracket 4, 7 \rrbracket$ . Using Fact 4.6, it is easy to see that  $\varphi^4(1|\beta_4|1) = \varphi^4(1|\gamma_4|1) = \varphi^4(\omega_1 1|\epsilon^1|1) = 0$ , whereas Fact 4.7 gives us immediately the identities  $\varphi^5(1|\alpha_4|1) = \varphi^5(1|\gamma_4|1) =$



$\varphi^5(\omega_1 1 | \epsilon^1 | 1) = 0$ , Fact 4.8 tells us that  $\varphi^6(1 | \alpha_4 | 1) = \varphi^6(1 | \beta_4 | 1) = \varphi^6(\omega_1 1 | \epsilon^1 | 1) = 0$ , and Fact 4.9 yields that  $\varphi^7(1 | \beta_4 | 1) = \varphi^7(1 | \gamma_4 | 1) = 0$  and  $\varphi^7(\omega_1 1 | \epsilon^1 | 1) = 3(bc - ba - ac)$ . For  $i \in \{4, 7\}$  and  $u \in \mathfrak{B}_4^{!*} \setminus \{\beta_4, \gamma_4\}$  (resp.,  $i = 5$  and  $u \in \mathfrak{B}_4^{!*} \setminus \{\alpha_4, \gamma_4\}$ ,  $i = 6$  and  $u \in \mathfrak{B}_4^{!*} \setminus \{\alpha_4, \beta_4\}$ ), we have that

$$\varphi^i(1 | u | 1) = (\rho^i Y_{14} - Y_{14} \rho_4^i)(1 | u | 1) = -Y_{14}(q_u^i) = -\lambda_i^u,$$

where  $\lambda_i^u \in \mathbb{k}$  is the coefficient of  $\omega_1 1 | \epsilon^1 | 1$  in  $q_u^i$ . It is easy to check that

$$\begin{aligned} \rho_3^4 \delta_4^b(1 | \alpha_4 | 1) &= bac | \alpha_3 | 1 + abc | \beta_3 | 1 + aba | \alpha_2 \gamma | 1 + v_{\alpha_4}^4, \\ \rho_3^4 \delta_4^b(1 | \alpha_3 \beta | 1) &= -2aba | \alpha_3 | 1 + 3abc | \gamma_3 | 1 + bac | \alpha_2 \beta | 1 + abc | \alpha_2 \gamma | 1 \\ &\quad + 2aba | \alpha \beta_2 | 1 + v_{\alpha_3 \beta}^4, \\ \rho_3^4 \delta_4^b(1 | \alpha_3 \gamma | 1) &= -2aba | \beta_3 | 1 + 2bac | \gamma_3 | 1 + 2aba | \alpha_2 \beta | 1 + bac | \alpha_2 \gamma | 1 \\ &\quad + abc | \alpha \beta_2 | 1 + v_{\alpha_3 \gamma}^4, \\ \rho_3^4 \delta_4^b(1 | \alpha_2 \beta_2 | 1) &= -2bac | \alpha_3 | 1 - 2abc | \beta_3 | 1 + 4aba | \gamma_3 | 1 + 2abc | \alpha_2 \beta | 1 \\ &\quad + 2bac | \alpha \beta_2 | 1 + v_{\alpha_2 \beta_2}^4, \end{aligned}$$

and

$$\begin{aligned} \rho_3^5 \delta_4^b(1 | \beta_4 | 1) &= bac | \alpha_3 | 1 + abc | \beta_3 | 1 + aba | \alpha_2 \gamma | 1 + v_{\beta_4}^5, \\ \rho_3^5 \delta_4^b(1 | \alpha_3 \beta | 1) &= -2aba | \alpha_3 | 1 + 2abc | \gamma_3 | 1 + bac | \alpha_2 \beta | 1 + aba | \alpha \beta_2 | 1 + v_{\alpha_3 \beta}^5, \\ \rho_3^5 \delta_4^b(1 | \alpha_3 \gamma | 1) &= abc | \alpha_3 | 1 - 2aba | \beta_3 | 1 + 2bac | \gamma_3 | 1 + 2aba | \alpha_2 \beta | 1 + bac | \alpha_2 \gamma | 1 \\ &\quad + abc | \alpha \beta_2 | 1 + v_{\alpha_3 \gamma}^5, \\ \rho_3^5 \delta_4^b(1 | \alpha_2 \beta_2 | 1) &= -3bac | \alpha_3 | 1 - 2abc | \beta_3 | 1 + 3aba | \gamma_3 | 1 + 2abc | \alpha_2 \beta | 1 \\ &\quad - aba | \alpha_2 \gamma | 1 + bac | \alpha \beta_2 | 1 + v_{\alpha_2 \beta_2}^5, \end{aligned}$$

as well as

$$\begin{aligned} \rho_3^6 \delta_4^b(1 | \gamma_4 | 1) &= -abc | \beta_3 | 1 + aba | \gamma_3 | 1 + bac | \alpha \beta_2 | 1 + v_{\gamma_4}^6, \\ \rho_3^6 \delta_4^b(1 | \alpha_3 \beta | 1) &= 4aba | \alpha_3 | 1 - 2bac | \beta_3 | 1 + 2abc | \alpha_2 \gamma | 1 + 2aba | \alpha \beta_2 | 1 + v_{\alpha_3 \beta}^6, \\ \rho_3^6 \delta_4^b(1 | \alpha_3 \gamma | 1) &= -abc | \alpha_3 | 1 + 2aba | \beta_3 | 1 + aba | \alpha_2 \beta | 1 + bac | \alpha_2 \gamma | 1 + v_{\alpha_3 \gamma}^6, \\ \rho_3^6 \delta_4^b(1 | \alpha_2 \beta_2 | 1) &= 4bac | \alpha_3 | 1 + 4abc | \beta_3 | 1 + 4aba | \alpha_2 \gamma | 1 + v_{\alpha_2 \beta_2}^6, \end{aligned}$$

together with

$$\begin{aligned} \rho_3^7 \delta_4^b(1 | \alpha_4 | 1) &= (aba - abc) | \alpha_3 | 1 + v_{\alpha_4}^7, \\ \rho_3^7 \delta_4^b(1 | \alpha_3 \beta | 1) &= -abc | \alpha_3 | 1 + 3bac | \alpha_3 | 1 + 3aba | \beta_3 | 1 + 2abc | \beta_3 | 1 - aba | \gamma_3 | 1 \\ &\quad - 2bac | \gamma_3 | 1 - abc | \alpha_2 \beta | 1 + 2aba | \alpha_2 \gamma | 1 + v_{\alpha_3 \beta}^7, \\ \rho_3^7 \delta_4^b(1 | \alpha_3 \gamma | 1) &= 2aba | \alpha_3 | 1 + 2bac | \alpha_3 | 1 + 2abc | \beta_3 | 1 - 2bac | \beta_3 | 1 - 2aba | \gamma_3 | 1 \\ &\quad - 2abc | \gamma_3 | 1 - abc | \alpha_2 \beta | 1 + aba | \alpha_2 \gamma | 1 + abc | \alpha_2 \gamma | 1 + aba | \alpha \beta_2 | 1 \end{aligned}$$

$$- bac|\alpha\beta_2|1 + v_{\alpha_3\gamma}^7,$$

$$\rho_3^7\delta_4^b(1|\alpha_2\beta_2|1) = aba|\alpha_3|1 - abc|\alpha_3|1 + aba|\beta_3|1 - abc|\gamma_3|1 + v_{\alpha_2\beta_2}^7,$$

where  $v_u^i \in \oplus_{j \in [0,2]} (A_j \otimes (A_{-3}^!)^* \otimes A_{3-j})$  if  $i \in \{4, 7\}$  and  $u \in \mathfrak{B}_4^{!*} \setminus \{\beta_4, \gamma_4\}$ , or if  $i = 5$  and  $u \in \mathfrak{B}_4^{!*} \setminus \{\alpha_4, \gamma_4\}$ , or if  $i = 6$  and  $u \in \mathfrak{B}_4^{!*} \setminus \{\alpha_4, \beta_4\}$ . Since,  $q_u^i$  is of the form  $q_u^i = B_u^i + \lambda_i^u \omega_1 | \epsilon^! | 1$  by degree reasons, where  $B_u^i \in K_4^b$ , and we have by definition that  $\delta_4^b(q_u^i) = \rho_3^i \delta_4^b(1|u|1)$ , we see that

$$d_4^b(B_u^i) = \delta_4^b(q_u^i) - \lambda_i^u f_0^b(1|\epsilon^!|1) = \rho_3^i \delta_4^b(1|u|1) - \lambda_i^u f_0^b(1|\epsilon^!|1) \tag{4.5}$$

for  $i \in [4, 7]$ . Using the explicit expression of the differential  $d_4^b$  given in [3], Fact 3.1, it is clear that the coefficients of  $aba|\gamma_3|1$  and  $aba|\alpha_2\gamma|1$  in  $d_4^b(B)$  coincide for all  $B \in K_4^b$ . Comparing the coefficients of  $aba|\gamma_3|1$  and  $aba|\alpha_2\gamma|1$  in both sides of the equation (4.5), together with the expression of  $f_0^b(1|\epsilon^!|1)$  given in [3], (3.2), we get

$$\lambda_4^{\alpha_4} = \lambda_5^{\beta_4} = -\lambda_6^{\gamma_4} = 1/3, \quad \lambda_i^{\alpha_3\beta} = \lambda_i^{\alpha_3\gamma} = 0, \quad \lambda_i^{\alpha_2\beta_2} = -\tau_i 4/3,$$

$$\lambda_7^{\alpha_4} = \lambda_7^{\alpha_2\beta_2} = 0, \quad \text{and} \quad \lambda_7^{\alpha_3\beta} = \lambda_7^{\alpha_3\gamma} = 1,$$

for  $i \in [4, 6]$ , where  $\tau_i = 1$  if  $i \in [4, 5]$  and  $\tau_6 = -1$ . Hence, we obtain that

$$\varphi^4(1|\alpha_4|1) = -1/3, \quad \varphi^4(1|\alpha_3\beta|1) = \varphi^4(1|\alpha_3\gamma|1) = 0, \quad \varphi^4(1|\alpha_2\beta_2|1) = 4/3,$$

$$\varphi^5(1|\beta_4|1) = -1/3, \quad \varphi^5(1|\alpha_3\beta|1) = \varphi^5(1|\alpha_3\gamma|1) = 0, \quad \varphi^5(1|\alpha_2\beta_2|1) = 4/3,$$

$$\varphi^6(1|\gamma_4|1) = 1/3, \quad \varphi^6(1|\alpha_3\beta|1) = \varphi^6(1|\alpha_3\gamma|1) = 0, \quad \varphi^6(1|\alpha_2\beta_2|1) = -4/3,$$

$$\varphi^7(1|\alpha_4|1) = \varphi^7(1|\alpha_2\beta_2|1) = 0 \quad \text{and} \quad \varphi^7(1|\alpha_3\beta|1) = \varphi^7(1|\alpha_3\gamma|1) = -1.$$

In consequence, we get

$$[-X_4, X_{14}] = (4/3)\alpha_2\beta_2|1 - (1/3)\alpha_4|1,$$

$$[-X_5, X_{14}] = (4/3)\alpha_2\beta_2|1 - (1/3)\beta_4|1,$$

$$[-X_6, X_{14}] = (1/3)\gamma_4|1 - (4/3)\alpha_2\beta_2|1,$$

$$[-X_7, X_{14}] = -(\alpha_3\beta + \alpha_3\gamma)|1 + 3\omega_1^* \epsilon^! |(bc - ba - ac).$$

Using the coboundaries  $g_{j,2}^2 \in \tilde{\mathfrak{B}}_2^2$  for  $j \in [4, 6]$  and  $e_{k,3}^3 \in \tilde{\mathfrak{B}}_3^3$  for  $k \in [7, 8]$  given in [3], Subsubsection 5.3.1, (4.3) as well as the identities

$$\alpha_2\beta|abc = X_4X_{10}, \quad \alpha_4|1 = X_9^2, \quad \text{and} \quad \alpha_2\beta_2|1 = X_9X_{10}, \tag{4.6}$$

which follow from [3], Fact 6.3 and (6.2), we can rewrite several brackets as

$$[-X_4, X_{10}] = \alpha\beta|(ab + bc) - \alpha\gamma|ac - g_{5,2}^2 = 0,$$

$$[-X_4, X_{11}] = -\alpha\beta|ab - \alpha\gamma|ba + g_{4,2}^2 = 0,$$

$$[-X_4, X_{12}] = \alpha_2|(bc - ba - ac) - g_{6,2}^2 = -2\alpha_2|(ab + ba) = -2X_1X_9,$$

$$\begin{aligned}[-X_4, X_{13}] &= 3\alpha_2\gamma|aba - 5\alpha\beta_2|bac - 5e_{7,3}^3 - 3e_{8,3}^3 = -8\alpha_2\beta|abc = -8X_4X_{10}, \\[-X_4, X_{14}] &= (4/3)\alpha_2\beta_2|1 - (1/3)\alpha_4|1 = (4/3)X_9X_{10} - (1/3)X_9^2.\end{aligned}$$

Analogously, using the coboundaries  $g_{j,2}^2 \in \tilde{\mathfrak{B}}_2^2$  for  $j \in \{4, 5, 7\}$  and  $e_{8,3}^3 \in \tilde{\mathfrak{B}}_3^3$  given in [3], Subsubsection 5.3.1, (4.3), (4.6) and the identity  $\beta_4|1 = X_{10}^2$  given in [3], Fact 6.3, we get that

$$\begin{aligned}[-X_5, X_9] &= -\alpha\beta|bc + \alpha\gamma|(ba + ac) - g_{4,2}^2 + g_{5,2}^2 = 0, \\[-X_5, X_{11}] &= -\alpha\beta|ab - \alpha\gamma|ba + g_{4,2}^2 = 0, \\[-X_5, X_{12}] &= -\beta_2|(ab + bc - ac) - g_{7,2}^2 = -2\beta_2|(ab + ba) = -2X_1X_{10}, \\[-X_5, X_{13}] &= -5\alpha_2\beta|abc + 3\alpha_2\gamma|aba - 3e_{8,3}^3 = -8\alpha_2\beta|abc = -8X_4X_{10}, \\[-X_5, X_{14}] &= (4/3)\alpha_2\beta_2|1 - (1/3)\beta_4|1 = (4/3)X_9X_{10} - (1/3)X_{10}^2.\end{aligned}$$

Moreover, using the coboundaries  $g_{j,2}^2 \in \tilde{\mathfrak{B}}_2^2$  for  $j \in \llbracket 1, 5 \rrbracket$  and  $e_{k,3}^3 \in \tilde{\mathfrak{B}}_3^3$  for  $k \in \llbracket 7, 8 \rrbracket$  given in [3], Subsubsection 5.3.1, (4.3), (4.6) and the identity  $\gamma_4|1 = X_{11}^2$  given in [3], Fact 6.3, we obtain

$$\begin{aligned}[-X_6, X_9] &= \alpha\beta|(bc - ab) - \alpha\gamma|(2ba + ac) + 2g_{4,2}^2 - g_{5,2}^2 = 0, \\[-X_6, X_{10}] &= -\alpha\beta|(ab + bc) + \alpha\gamma|ac + g_{5,2}^2 = 0, \\[-X_6, X_{12}] &= \gamma_2|(bc - ba - ac) - \alpha\beta|ba - \alpha\gamma|ab - 2g_{1,2}^2 - 2g_{2,2}^2 - g_{3,2}^2 \\ &= -2(\alpha_2 + \beta_2)|(ab + ba) = -2X_1(X_9 + X_{10}), \\[-X_6, X_{13}] &= -10\alpha_2\gamma|aba - 2\alpha\beta_2|bac - 2e_{7,3}^3 + 10e_{8,3}^3 = 8\alpha_2\beta|abc = 8X_4X_{10}, \\[-X_6, X_{14}] &= (1/3)\gamma_4|1 - (4/3)\alpha_2\beta_2|1 = (1/3)X_{11}^2 - (4/3)X_9X_{10}.\end{aligned}$$

Finally, using the coboundaries  $g_{j,2}^2 \in \tilde{\mathfrak{B}}_2^2$  for  $j \in \llbracket 1, 6 \rrbracket \setminus \{3\}$  and  $e_{k,3}^3 \in \tilde{\mathfrak{B}}_3^3$  for  $k \in \llbracket 1, 4 \rrbracket \cup \llbracket 9, 10 \rrbracket$  given in [3], Subsubsection 5.3.1, (4.3) and

$$\begin{aligned}\alpha_3|(aba - abc) &= X_7X_9, & \alpha_3|aba + \beta_3|bac &= X_7(X_9 + X_{10}) - 2X_6X_{12}, \\(\alpha_3 + \beta_3)|aba &= X_6X_{12}, & (\alpha_3\beta + \alpha_3\gamma)|1 + 3\omega_1\epsilon^1|(ba - bc + ac) &= X_9X_{12},\end{aligned}$$

given in [3], Fact 6.3, or in [3], (6.2), together with the second element in the fifth and the eighth line, the first element in the ninth line of [3] (6.5), we have that

$$\begin{aligned}[-X_7, X_9] &= \alpha_2|(bc - ab - ac - 2ba) - \alpha\beta|(ba + ac) + \alpha\gamma|bc - g_{1,2}^2 - g_{6,2}^2 \\ &= -4\alpha_2|(ab + ba) = -4X_1X_9, \\[-X_7, X_{10}] &= \alpha\beta|ac - \alpha\gamma|(ab + bc) - g_{2,2}^2 = -\beta_2|(ab + ba) = -X_1X_{10}, \\[-X_7, X_{11}] &= \alpha\beta|ba + \alpha\gamma|ab + g_{1,2}^2 + g_{2,2}^2 = (\alpha_2 + \beta_2)|(ab + ba) \\ &= X_1(X_9 + X_{10}), \\[-X_7, X_{12}] &= (\alpha\beta + \alpha\gamma)|(bc - ba - ac) - g_{1,2}^2 + g_{4,2}^2 - g_{5,2}^2 = -\alpha_2|(ab + ba)\end{aligned}$$

$$\begin{aligned}
 &= -X_1X_9, \\
 [-X_7, X_{13}] &= \alpha_3|(abc - 2aba) + \alpha_2\beta|(2bac - 6aba) - \alpha_2\gamma|(abc + 4bac) \\
 &\quad + 5\alpha\beta_2|(abc - aba) - (1/3)(23e_{1,3}^3 + 11e_{2,3}^3 - 32e_{3,3}^3 - 16e_{4,3}^3 \\
 &\quad - 5e_{9,3}^3 + 6e_{10,3}^3) \\
 &= (8/3)\alpha_3|(aba - abc) - (32/3)(\alpha_3 + \beta_3)|aba \\
 &\quad - (16/3)(\alpha_3|aba + \beta_3|bac) \\
 &= -(8/3)X_7(X_9 + 2X_{10}) = -4X_1X_{13} + 4X_2X_{13} + 8X_4X_{12}, \\
 [-X_7, X_{14}] &= -(\alpha_3\beta + \alpha_3\gamma)|1 + 3\omega_1^*\epsilon^1|(bc - ba - ac) = -X_9X_{12}.
 \end{aligned}$$

The proposition is thus proved. □

REMARK 4.11. *Note that vanishing of  $[X_i, X_j]$  for  $i \in \llbracket 4, 7 \rrbracket$  and  $j \in \llbracket 3, 7 \rrbracket$  in Proposition 4.10 also follows from a simple degree argument based on Corollary 2.5 and the Hilbert series of the Hochschild cohomology given in [3], Cor. 5.9.*

### 4.3 GERSTENHABER BRACKETS

We will finally compute the remaining Gerstenhaber brackets. We start with the following result, which is a sort of descending argument.

LEMMA 4.12. *Let  $H = \bigoplus_{n \in \mathbb{N}_0} H^n$  be a Gerstenhaber algebra with bracket  $[\cdot, \cdot]$ . Let  $x \in H^{n+1}$ ,  $y \in H^n$ ,  $a_x \in H^0$ ,  $a_y \in H^1$  and  $z \in H^m$  satisfy that  $a_x x = a_y y$ , and there is a vector subspace  $M \subseteq H^{n+m-1}$  such that  $[y, z] \in M$  and the map  $\mu_{a_y} : M \rightarrow H^{n+m}$  sending  $v \in M$  to  $a_y v$  is injective. Then,  $[y, z]$  is the unique element  $v \in M$  satisfying that  $a_y v$  coincides with*

$$(-1)^{m-1}(a_x[x, z] + [a_x, z]x - [a_y, z]y). \tag{4.7}$$

*Proof.* By (2.2) we get that

$$[a_x x, z] = [a_x, z]x + a_x[x, z] \text{ and } [a_y y, z] = [a_y, z]y + (-1)^{m-1}a_y[y, z].$$

These identities together with  $a_x x = a_y y$  imply

$$a_y[y, z] = (-1)^{m-1}(a_x[x, z] + [a_x, z]x - [a_y, z]y).$$

Hence, the right member is in the image of the injective map  $\mu_{a_y}$ , and the result follows. □

REMARK 4.13. *We will apply the previous lemma to the case when  $H = \text{HH}^\bullet(A)$  is the Hochschild cohomology of a graded algebra  $A$ , so  $H$  is endowed with an extra grading, called internal (see Corollary 2.5), the elements  $x, y, z, a_x, a_y$  are homogeneous for both gradings and  $M \subseteq H^{n+m-1}$  is the subspace of internal degree equal to the sum of those of  $y$  and  $z$ . In this case, the methods given in Subsections 2.2 and 2.3 allow to compute the last two brackets of (4.7), whereas the first one will usually vanish by degree reasons.*

PROPOSITION 4.14. *Let  $A = \mathrm{FK}(3)$  be the Fomin-Kirillov algebra on 3 generators. Then, we have the Gerstenhaber brackets  $[X_i, X_j] = 0$  for  $i, j \in \llbracket 9, 14 \rrbracket \setminus \{13\}$  and*

$$[X_{13}, X_j] = \begin{cases} 2X_j^2, & \text{if } j \in \llbracket 9, 11 \rrbracket, \\ -6X_1X_{14} + 6X_2X_{14} + 2X_9X_{12}, & \text{if } j = 12, \\ 0, & \text{if } j = 13, \\ 4(X_9 + X_{10} + X_{11})X_{14}, & \text{if } j = 14. \end{cases}$$

*Proof.* Recall that, by Corollary 2.5, the Gerstenhaber bracket satisfies that  $[,] : H_{m_1}^{n_1} \times H_{m_2}^{n_2} \rightarrow H_{m_1+m_2-1}^{n_1+n_2-1}$ , where  $H_{m_i}^{n_i}$  has internal degree  $m_i - n_i$  for  $i = 1, 2$ . Using this degree argument together with the Hilbert series of the Hochschild cohomology computed in [3], Cor. 5.9, we easily see that  $[X_i, X_j] = 0$  for  $i, j \in \llbracket 9, 14 \rrbracket \setminus \{13\}$ . Moreover,  $[X_{13}, X_{13}] = 0$  by (2.1).

It remains to compute  $[X_{13}, X_j]$  for all  $j \in \llbracket 9, 14 \rrbracket \setminus \{13\}$ . Note first the identities

$$\begin{aligned} [X_8, X_9]X_{13} - 6[X_3, X_9]X_{14} &= 2X_9X_{13} + 12X_4X_{14} = 2X_8X_9^2, \\ [X_8, X_{10}]X_{13} - 6[X_3, X_{10}]X_{14} &= 2X_{10}X_{13} + 12X_5X_{14} = 2X_8X_{10}^2, \\ [X_8, X_{11}]X_{13} - 6[X_3, X_{11}]X_{14} &= 2X_{11}X_{13} - 12X_6X_{14} = 2X_8X_{11}^2, \\ [X_8, X_{12}]X_{13} - 6[X_3, X_{12}]X_{14} &= 2X_{12}X_{13} - 12X_7X_{14} + 6X_1X_8X_{14} \\ &\quad - 6X_2X_8X_{14} \\ &= 2X_8X_{11}X_{12} + 6X_1X_8X_{14} \\ &= 2X_8X_{12}^2 - 4X_8X_9X_{10} \\ &= 2X_8X_9X_{12} - 6X_1X_8X_{14} + 6X_2X_8X_{14}, \\ [X_8, X_{14}]X_{13} - 6[X_3, X_{14}]X_{14} &= 4X_8(X_9 + X_{10} + X_{11})X_{14}, \end{aligned} \tag{4.8}$$

where the first equality of the first four lines as well as that of the last line follows from Propositions 4.4 and 4.5, and we used the first element of the seventh and the eighth line of [3], (6.5), as well as its last four elements, for the remaining equalities. The penultimate element of the ninth line of [3], (6.5), also tells us that  $6X_3X_{14} = X_8X_{13} \in \mathrm{HH}^\bullet(A)$ .

Notice now that, by degree reasons,  $[X_{13}, X_j] \in H_0^4$  for  $j \in \llbracket 9, 12 \rrbracket$  and  $H_0^4$  is precisely the subspace of  $\mathrm{HH}^4(A)$  spanned by the elements  $X_9^2, X_{10}^2, X_{11}^2, X_9X_{12} - 3X_1X_{14} + 3X_2X_{14}, X_9X_{10}, X_1X_{14}$  and  $X_2X_{14}$ . On the other hand,  $[X_{13}, X_{14}] \in H_{-2}^6 = \omega_1^*H_0^2$ , by degree reasons, and  $\omega_1^*H_0^2$  is the subspace of  $\mathrm{HH}^4(A)$  spanned by  $X_9X_{14}, X_{10}X_{14}, X_{11}X_{14}$  and  $X_{12}X_{14}$ . Let us denote by  ${}^jM \subseteq \mathrm{HH}^4(A)$  the subspace given by  $H_0^4$  if  $j \in \llbracket 9, 12 \rrbracket$  and by  $H_{-2}^6$  if  $j = 14$ . Since the elements  $X_8X_9^2, X_8X_{10}^2, X_8X_{11}^2, X_8X_9X_{12} - 3X_1X_8X_{14} + 3X_2X_8X_{14}, X_8X_9X_{10}, X_1X_8X_{14}$ , and  $X_2X_8X_{14}$  are linearly independent, by the second equalities of the first four lines of (4.8) together with [3], (6.7) and (6.8), the map  ${}^jM \rightarrow \mathrm{HH}^5(A)$  given by left multiplication by  $X_8$  is injective for  $j \in \llbracket 9, 12 \rrbracket$ . Similarly, the elements  $X_8X_9X_{14}, X_8X_{10}X_{14}, X_8X_{11}X_{14}$  and  $X_8X_{12}X_{14}$  are

linearly independent, by [3], (6.8), so the map  ${}^{14}M \rightarrow \text{HH}^7(A)$  given by left multiplication by  $X_8$  is also injective.

Finally, applying Lemma 4.12 to  $x = X_{14}$ ,  $y = X_{13}$ ,  $z = X_j$ ,  $a_x = 6X_3$ ,  $a_y = X_8$  and  $M = {}^jM$  for  $j \in \llbracket 9, 14 \rrbracket \setminus \{13\}$ , together with the fact remarked at the beginning of the proof that  $[X_{14}, X_j] = 0$  and (4.8), the result follows.  $\square$

We can summarize the calculations of the Gerstenhaber brackets on  $\text{HH}^\bullet(A)$  done in Propositions 4.4, 4.5, 4.10 and 4.14 in the following table, where the brackets strictly below the diagonal are not displayed since they can be obtained using Lemma 2.2.

$\rho \setminus \phi$	$X_1$	$X_2$	$X_3$	$X_4$	$X_5$	$X_6$	$X_7$	$X_8$	$X_9$	$X_{10}$	$X_{11}$	$X_{12}$	$X_{13}$	$X_{14}$
$X_1$	0	0	0	0	0	0	0	$2X_1$	0	0	0	0	$4X_1(X_9 + X_{10})$	0
$X_2$	0	0	0	0	0	0	0	$2X_2$	0	0	0	0	$4X_1X_{10}$	0
$X_3$		0	0	0	0	0	0	$4X_3$	$-2X_4$	$-2X_5$	$2X_6$	$2X_7 - X_1X_8 + X_2X_8$	$4X_3(X_9 + X_{10} + X_{11})$	$X_{13} - (2/3)X_8(X_9 + X_{10} + X_{11})$
$X_4$			0	0	0	0	$2X_1$	0	0	0	0	$2X_1X_9$	$8X_4X_{10}$	$(1/3)X_8^2 - (4/3)X_9X_{10}$
$X_5$				0	0	0	$2X_5$	0	0	0	0	$2X_1X_{10}$	$8X_4X_{10}$	$(1/3)X_{10}^2 - (4/3)X_9X_{10}$
$X_6$					0	0	$2X_6$	0	0	0	0	$2X_1(X_9 + X_{10})$	$-8X_4X_{10}$	$(4/3)X_9X_{10} - (1/3)X_8^2$
$X_7$						0	$2X_7$	$4X_1X_9$	$X_1X_{10}$	$-X_1(X_9 + X_{10})$	$X_1X_9$	$4X_1X_{13} - 4X_2X_{13} - 8X_4X_{12}$	$2X_{13}$	$X_9X_{12}$
$X_8$							0	$2X_9$	$2X_{10}$	$2X_{11}$	$2X_{12}$	$2X_{13}$	$2X_{12}$	$6X_{14}$
$X_9$								0	0	0	0	$-2X_9^2$	0	0
$X_{10}$									0	0	0	$-2X_{10}^2$	0	0
$X_{11}$										0	0	$-2X_{11}^2$	0	0
$X_{12}$											0	0	$6X_1X_{14} - 6X_2X_{14} - 2X_9X_{12}$	0
$X_{13}$												0	0	$4(X_9 + X_{10} + X_{11})X_{14}$
$X_{14}$														0

Table 4.1: Gerstenhaber brackets  $[\rho, \phi]$ .

PROPOSITION 4.15. *There is no generator of the Gerstenhaber bracket on the Hochschild cohomology  $\text{HH}^\bullet(A)$  of  $A = \text{FK}(3)$ , i.e. there is no map  $\Delta : \text{HH}^\bullet(A) \rightarrow \text{HH}^\bullet(A)$  of degree  $-1$  such that*

$$[x, y] = (-1)^{|x|}(\Delta(xy) - \Delta(x)y - (-1)^{|x|}x\Delta(y)) \tag{4.9}$$

for all homogeneous elements  $x, y \in \text{HH}^\bullet(A)$ , where  $|x|$  is the cohomological degree of  $x$ . In particular, there is no Batalin-Vilkovisky structure on  $\text{HH}^\bullet(A)$  inducing the Gerstenhaber bracket.

*Proof.* Assume that (4.9) holds. Obviously,  $\Delta(\text{HH}^0(A)) = 0$ . Applying the results in Table 4.1 and (4.9), we get  $-4X_3 = [X_8, X_3] = \Delta(X_8)X_3$ , and  $0 = [X_i, X_j] = \Delta(X_i)X_j$  for  $i \in \llbracket 4, 7 \rrbracket$  and  $j \in \llbracket 1, 3 \rrbracket$ , since  $X_8X_3 = X_iX_j = 0$  in that case (see the first two lines of [3], (6.5)). Hence,  $\Delta(X_8) \in -4 + \text{span}_{\mathbb{k}}\langle X_1, X_2, X_3 \rangle$  and  $\Delta(X_i) \in \text{span}_{\mathbb{k}}\langle X_1, X_2, X_3 \rangle$  for  $i \in \llbracket 4, 7 \rrbracket$ , where  $\text{span}_{\mathbb{k}}\langle X_1, X_2, X_3 \rangle$  is the  $\mathbb{k}$ -subspace spanned by  $X_1, X_2, X_3$ . Moreover,

$$\begin{aligned} -2X_4 &= [X_3, X_9] = \Delta(X_3X_9) - X_3\Delta(X_9) = \Delta(X_3X_9), \\ 2X_4 &= [X_4, X_8] = -\Delta(X_4X_8) + \Delta(X_4)X_8 - X_4\Delta(X_8) \\ &= -\Delta(X_4X_8) + \Delta(X_4)X_8 + 4X_4, \end{aligned} \tag{4.10}$$

where we used that  $X_4X_i = X_3X_k = 0$  for  $i \in \llbracket 1, 3 \rrbracket$  and  $k \in \llbracket 4, 8 \rrbracket$ , by the first two lines of [3], (6.5). Since  $X_3X_9 = X_4X_8 \in \text{HH}^2(A)$  (see the

penultimate element of the third line of [3], (6.5)), adding the equations (4.10), we obtain  $\Delta(X_4)X_8 + 4X_4 = 0$ . The identity  $\Delta(X_4) = k_1X_1 + k_2X_2 + k_3X_3$  for  $k_1, k_2, k_3 \in \mathbb{k}$ , which we proved before, implies that  $k_1X_1X_8 + k_2X_2X_8 + 4X_4 = 0$ . This is impossible since the elements  $X_1X_8, X_2X_8$  and  $X_4$  are linearly independent in  $\mathrm{HH}^1(A)$  (see [3], (6.7)). The proposition thus follows.  $\square$

## REFERENCES

- [1] Sergey Fomin and Anatol N. Kirillov, *Quadratic algebras, Dunkl elements, and Schubert calculus*, Advances in geometry, Progr. Math., 172. Birkhäuser Boston, Boston, MA, 1999, pp. 147–182. MR1667680 ↑1782, 1783
- [2] Murray Gerstenhaber, *The cohomology structure of an associative ring*, Ann. of Math. (2) 78 (1963), 267–288, DOI 10.2307/1970343. MR161898 ↑1776
- [3] Estanislao Herscovich and Ziling Li, *Hochschild and cyclic (co)homology of the Fomin–Kirillov algebra on 3 generators* (2021), available at <https://arxiv.org/abs/2112.01201>. ↑1773, 1774, 1783, 1784, 1785, 1789, 1790, 1798, 1799, 1800, 1801, 1802, 1803
- [4] Alexander Milinski and Hans-Jürgen Schneider, *Pointed indecomposable Hopf algebras over Coxeter groups*, New trends in Hopf algebra theory (La Falda, 1999), Contemp. Math., 267. Amer. Math. Soc., Providence, RI, 2000, pp. 215–236, DOI 10.1090/conm/267/04272. MR1800714 ↑1782
- [5] Constantin Năstăsescu and Freddy Van Oystaeyen, *Methods of graded rings*, Lecture Notes in Mathematics, 1836. Springer, Berlin, 2004. MR2046303 ↑1777
- [6] Cris Negron and Sarah Witherspoon, *An alternate approach to the Lie bracket on Hochschild cohomology*, Homology Homotopy Appl. 18 (2016), no. 1, 265–285, DOI 10.4310/HHA.2016.v18.n1.a14. MR3498646 ↑1781
- [7] Dragoş Ştefan and Cristian Vay, *The cohomology ring of the 12-dimensional Fomin–Kirillov algebra*, Adv. Math. 291 (2016), 584–620, DOI 10.1016/j.aim.2016.01.001. MR3459024 ↑1783
- [8] Mariano Suárez-Álvarez, *A little bit of extra functoriality for Ext and the computation of the Gerstenhaber bracket*, J. Pure Appl. Algebra 221 (2017), no. 8, 1981–1998, DOI 10.1016/j.jpaa.2016.10.015. MR3623179 ↑1773, 1774, 1781, 1782, 1791
- [9] Yury Volkov, *Gerstenhaber bracket on the Hochschild cohomology via an arbitrary resolution*, Proc. Edinb. Math. Soc. (2) 62 (2019), no. 3, 817–836, DOI 10.1017/s0013091518000901. MR3974969 ↑1780

- [10] Sarah J. Witherspoon, *Hochschild cohomology for algebras*, Graduate Studies in Mathematics, 204. American Mathematical Society, Providence, RI, 2019. MR3971234 ↑1775, 1776

Estanislao Herscovich  
Institut Fourier  
UMR 5582  
Laboratoire de Mathématiques  
Université Grenoble Alpes  
CS 40700  
38058, Grenoble cedex 9  
France

Ziling Li  
Institut Fourier  
UMR 5582  
Laboratoire de Mathématiques  
Université Grenoble Alpes  
CS 40700  
38058, Grenoble cedex 9  
France